arXiv:0806.0281v1 [math.CO] 2 Jun 2008

k-flaw Preference Sets

Po-Yi Huang[a,*] Jun Ma[b,†] Jean Yeh[c,‡]

[a] Department of Mathematics, National Cheng Kung University, Tainan, Taiwan

[b] Institute of Mathematics, Academia Sinica, Taipei, Taiwan

[c] Department of Mathematics, National Taiwan University, Taipei, Taiwan

Abstract

In this paper, let $\mathcal{P}^{l}_{n;\leq s;k}$ denote a set of k-flaw preference sets $(a_1,\ldots,a_n)$ with n parking spaces satisfying that $1 \leq a_i \leq s$ for any i and $a_1 = l$ and $p^{l}_{n;\leq s;k} = |\mathcal{P}^{l}_{n;\leq s;k}|$. We use a combinatorial approach to the enumeration of k-flaw preference sets by their leading terms. The approach relies on bijections between the k-flaw preference sets and labeled rooted forests. Some bijective results between certain sets of k-flaw preference sets of distinct leading terms are also given. We derive some formulas and recurrence relations for the sequences $p^{l}_{n;\leq s;k}$ and give the generating functions for these sequences.

Keyword: Flaw; Leading term; Parking function; Forest

---

[*] Partially supported by NSC 96-2115-M-006-012

[†] Email address of the corresponding author: majun@math.sinica.edu.tw

[‡] jean.yh@ms45.url.com.tw

1 Introduction

Throughout the paper, let $[n] := \{1, 2, \ldots, n\}$ and $[m, n] := \{m, \ldots, n\}$. Suppose that n cars have to be parked in m parking spaces which are arranged in a line and numbered 1 to m from left to right. Each car has initial parking preference a_i; if space a_i is occupied, the car moves to the first unoccupied space to the right. We call $(a_1, \ldots, a_n)$ *preference set.* Clearly, the number of preference sets is m^n. If a preference set $(a_1, \ldots, a_n)$ satisfies $a_i \leq a_{i+1}$ for $1 \leq i \leq n-1$, then we say that the preference set is *ordered.* If all the cars can find a parking space, then we say the preference set is a *parking function.* If there are exactly k cars which can't be parked, then the preference set is called a *k-flaw preference set.*

Let n, m, s, and k be four nonnegative integers with $1 \leq s \leq m$ and $k \leq n-1$. Suppose that there are m parking spaces. We use $\mathcal{P}_{n,m;\leq s;k}$ to denote a set of k-flaw preference sets $(a_1, \ldots, a_n)$ of length n satisfying $1 \leq a_i \leq s$ for all i. For $1 \leq l \leq s$, we use $\mathcal{P}^l_{n,m;\leq s;k}$ to denote a set of preference sets $(a_1, \ldots, a_n) \in \mathcal{P}_{n,m;\leq s;k}$ such that $a_1 = l$. Let $p_{n,m;\leq s;k} = |\mathcal{P}_{n,m;\leq s;k}|$ and $p^l_{n,m;\leq s;k} = |\mathcal{P}^l_{n,m;\leq s;k}|$. For any of the above cases, if the parameter k (resp. m) doesn't appear, we understand $k = 0$ (resp. $m = n$); if the parameter m and s are both erased, we understand $s = m = n$.

There are some results about parking functions with $s = m = n$. Riordan introduced parking functions in [15]. He derived that the number of parking functions of length n is $(n+1)^{n-1}$, which coincides with the number of labeled trees on $n+1$ vertices by Cayley's formula. Several bijections between the two sets are known (e.g., see [7, 15, 16]). Furthermore, define a generating function $P(x) = \sum\limits_{n\geq 0} \frac{(n+1)^{n-1}}{n!} x^n$. It is well known that $xP(x)$ is the compositional inverse of the function $\psi(x) = xe^{-x}$, i.e., $\psi(xP(x)) = x$. Riordan concluded that the number of ordered parking functions is $\frac{1}{n+1}\binom{2n}{n}$, which is also equals the number of Dyck path of semilength n. Parking functions have been found in connection to many other combinatorial structures such as acyclic mappings, polytopes, non-crossing partitions, non-nesting partitions, hyperplane arrangements,etc. Refer to [6, 7, 8, 13, 17, 18] for more information.

Any parking function $(a_1, \ldots, a_n)$ can be redefined such that its increasing rearrangement $(b_1, \ldots, b_n)$ satisfies $b_i \leq i$. Pitman and Stanley generalized the notion of parking functions in

[13]. Let $\mathbf{x} = (x_1, \ldots, x_n)$ be a sequence of positive integers. The sequence $\alpha = (a_1, \ldots, a_n)$ is called an $\mathbf{x}$-parking function if the non-decreasing rearrangement $(b_1, \ldots, b_n)$ of α satisfies $b_i \leq x_1 + \ldots + x_i$ for any $1 \leq i \leq n$. Thus, the ordinary parking function is the case $\mathbf{x} = (1, \ldots, 1)$. By the determinant formula of Gončarove polynomials, Kung and Yan [12] obtained the number of $\mathbf{x}$-parking functions for an arbitrary $\mathbf{x}$. See also [19, 20, 21] for the explicit formulas and properties for some specified cases of $\mathbf{x}$.

An $\mathbf{x}$-parking function $(a_1, \ldots, a_n)$ is said to be k-leading if $a_1 = k$. Let $q_{n,k}$ denote the number of k-leading ordinary parking functions of length n. Foata and Riordan [7] derived a generating function for $q_{n,k}$ algebraically. Recently, Sen-peng Eu, Tung-shan Fu and Chun-Ju Lai [3] gave a combinatorial approach to the enumeration of $(a, b, \ldots, b)$-parking functions by their leading terms.

Riordan [15] told us the relations between ordered parking functions and Dyck paths. Sen-peng Eu et al. [4, 5] considered the problem of the enumerations of lattice paths with flaws. It is natural to consider the problem of the enumerations of preference sets with flaws. There is a interesting facts. Salmon and Cayley in 1849 established the classical configuration of 27 lines in a general cubic surface. Given a line l, the number of lines, which are disjoint from, intersect or are equal to l, are 16,10 and 1, respectively, see [9] for the detail information. These number exactly are the number of 0-, 1- and 2-flaw preference sets of length 3. Peter J Cameron et al. [1] indicate that there are some relations between k-flaw preference sets and the context of hashing since data would be lost. Also they counted the number of k-flaw preference sets and calculate the asymptotic. Ordered k-flaw preference sets were studied in [10] and some enumerations for some parking functions were given in [11].

In this paper, we use the methods developed by Sen-peng Eu et al. [3] to study k-flaw preference sets. Sen-peng Eu et al. find that *triplet-labelled rooted forests* enable $(a, 1, \ldots, 1)$-parking functions to be manipulated on forests easily. The methods are different with that in [1]. We consider triplet-labelled rooted forests associated with k-flaw preference sets and establish a bijection between k-flaw preference sets and label rooted forests, so as to enumerate k-flaw preference sets by leading term.

First, we enumerate k-flaw preference sets in the set $\mathcal{P}_{n;\leq s;k}$. Then we consider the triplet-labelled rooted forests and bijections associated with the parking functions in the set $\mathcal{P}_{n,n+k;\leq n+k}$. Using these bijections, we find that $p^1_{n,n+k;\leq n+k} = p^2_{n,n+k;\leq n+k} = \ldots = p^{k+1}_{n,n+k;\leq n+k}$ for any $k \geq 0$. Taking $l = 1$, we have (1) $p^1_{n,n+k;\leq n+k} = p_{n-1,n+k;\leq n+k}$ for any $k \geq 0$ and $n \geq 1$; (2) $p^1_{n,n+k,\leq n+k} = \sum\limits_{i=1}^{n} \binom{n-1}{i-1} p^1_{i,i+k-1,\leq i+k-1} p_{n-i}$ for any $k \geq 1$. When $k+1 \leq l \leq n+k-1$, we obtain the recurrence relation $p^l_{n,n+k;\leq n+k} - p^{l+1}_{n,n+k;\leq n+k} = \binom{n-1}{l-k-1} p_{l-k-1,l-1;\leq l-1} p_{n+k-l}$.

To enumerate k-flaw preference sets in the set $\mathcal{P}^l_{n;\leq s;k}$, we study the triplet-labelled rooted forests and bijections associated with the k-flaw preference sets in the set $\mathcal{P}_{n;\leq s;k}$. We prove that $p^1_{n;\leq s;k} = p^2_{n;\leq s;k} = \ldots = p^k_{n;\leq s;k}$ for any $k \geq 1$; letting $l = 1$, we have $p^1_{n;\leq s;k} = \sum\limits_{i=1}^{s-k-1} \binom{n-1}{s-i-k-1} p_{n+k-s+i;\leq i} p^1_{s-k-i,s-i-1;\leq s-i-1}$ for any $k \geq 1$ and $k+1 \leq s \leq n$. For the cases with $s = l$, we derive some interesting identities: (1) $p^s_{n;\leq s;k+1} = p^1_{n;\leq s;k}$ for any $k \geq 1$ and $k+2 \leq s \leq n$; (2) $p^s_{n+1;\leq s;k} = p_{n;\leq s;k}$ for $k \geq 0$ and $k+1 \leq s \leq n$; (3) $p^s_{n+1;\leq s;k} = p_{n;\leq s;k}$ for any $k \geq 0$ and $k+1 \leq s \leq n$; (4) $p^n_{n;1} = p^2_n$ for any $n \geq 2$; (5) $p^n_{n;1} - p^{n-1}_{n;1} = p_{n-2}$ for any $n \geq 2$. When $k \leq l \leq s-1$, we obtain the following recurrence relation

$$\begin{aligned} p^{l+1}_{n;\leq s;k} - p^l_{n;\leq s;k} \quad = \quad & \sum_{i=k-1}^{l-2} \binom{n-1}{i-k+1} p_{i-k+1,i;\leq i} \left[p^{l-i}_{n+k-i-1;\leq s-i-1} - p^{l-i-1}_{n+k-i-1;\leq s-i-1}\right] \\ & + \binom{n-1}{l-k} p_{l-k,l-1;\leq l-1} \left[p^1_{n+k-l;\leq s-l} - p^{s-l}_{n+k-l;\leq s-l;1}\right]. \end{aligned}$$

We also are interested in generating functions for some sequences. For any $k \geq 1$ and $s \geq 0$, define a generating function $D_{k,s}(x) = \sum\limits_{n \geq s+k} \frac{p_{n;\leq n-s;k}}{n!} x^n$, then $D_{k,s}(x) = [P(x)]^{k+1} \sum\limits_{i=0}^{k+s} \frac{(-1)^i (k+s+1-i)^i}{i!} x^i - [P(x)]^k \sum\limits_{i=0}^{k+s-1} \frac{(-1)^i (k+s-i)^i}{i!} x^i$. Furthermore, let $D(x,y,z) = \sum\limits_{k \geq 1} \sum\limits_{s \geq 0} \sum\limits_{n \geq s+k} \frac{p_{n;\leq n-s;k}}{n!} x^n y^s z^k$, then $D(x,y,z) = \frac{zP(x)}{y-zP(x)} \left[\frac{P(x)-y}{e^{xy}-y} - \frac{(1-z)P(x)}{e^{xzP(x)}-zP(x)}\right]$.

For any $l \geq 0$ and $k \geq 0$, define a generating function $H_{l,k}(x) = \sum\limits_{n \geq l+1} \frac{p^{n+k-l}_{n,n+k;\leq n+k}}{(n-1)!} x^n$, then $H_{l,k}(x)$ satisfies the recurrence relation $H_{l,k}(x) = H_{l-1,k}(x) - \frac{p^1_{l,l+k;\leq l+k}}{(l-1)!} x^l + \frac{p_l}{l!} x^{l+1} [P(x)]^{k+1}$ with the initial conditions $H_{0,k}(x) = x[P(x)]^{k+1}$. Let $H_k(x,y) = \sum\limits_{l \geq 0} H_{l,k}(x) y^l$ and $H(x,y,z) = \sum\limits_{k \geq 0} H_k(x,y) z^k$, then $H_k(x,y) = \frac{xP(xy)\{[P(x)]^{k+1} - y[P(xy)]^{k+1}\}}{1-y}$ and $H(x,y,z) = \frac{xP(xy)}{1-y} \left[\frac{P(x)}{1-zP(x)} - \frac{yP(xy)}{1-zP(xy)}\right]$.

Define a generating function $W(x,y,z,v)=\sum\limits_{k\geq 1}\sum\limits_{s\geq 0}\sum\limits_{l\geq s}\sum\limits_{n\geq k+l}\frac{p^{n-l}_{n;\leq n-s;k}}{(n-1)!}x^ny^lz^sv^k$, then

$$\begin{aligned}W(x,y,z,v) &= \frac{xyvP(xy)}{y-1}\left\{\left[\frac{yP(xy)}{yz-vP(xy)}-\frac{P(x)}{yz-vP(x)}\right]R(xy,z)\right.\\ &\quad\left.+\frac{P(x)}{yz-vP(x)}R(xy,\frac{v}{y}P(x))-\frac{yP(xy)}{yz-vP(xy)}R(xy,\frac{v}{y}P(xy))\right\}\\ &\quad+\frac{vP(x)}{yz-vP(x)}\left[F(x,y,z)-F(x,y,\frac{v}{y}P(x))\right]\end{aligned}$$

where $R(x,y)=\frac{P(x)-y}{e^{xy}-y}$ and $F(x,y,z)=\frac{x}{e^{xyz}-z}\left[\frac{P(xy)(P(x)-yP(xy))}{1-y}-\frac{zP(xyz)[P(x)-yzP(xyz)]}{1-yz}\right]$.

Recently, Postnikov and Shapiro [14] gave a new generalization, building on work of Cori, Rossin and Salvy [2], the G-parking functions of a graph. For the complete graph $G=K_{n+1}$, the defined functions in [14] are exactly the classical parking functions. So, in the future work, we will consider k-flaw G-parking function.

We organize this paper as follows. In Section 2, we enumerate k-flaw preference sets in the set $\mathcal{P}_{n;\leq s;k}$. In Section 3, we consider the triplet-labelled rooted forests and bijections associated with the parking functions in the set $\mathcal{P}_{n,n+k;\leq n+k}$. In Section 4, we give the enumerations of parking functions in the set $\mathcal{P}^l_{n,n+k;\leq n+k}$. In Section 5, we study the triplet-labelled rooted forests and bijections associated with the k-flaw preference sets in the set $\mathcal{P}_{n;\leq s;k}$. In Section 6, we investigate the problems of the enumerations of preference sets in the set $\mathcal{P}^l_{n;\leq s;k}$. In Section 7, we obtain some generating functions for some sequences given in the previous sections. In Appendix, we list the values of $p^l_{n;\leq s;k}$ for $n\leq 7$ and $p^l_{n,n+k;\leq n+k}$ for any $n\leq 5$ and $k\leq 3$.

2 Counting the number of elements in $\mathcal{P}_{n;\leq s;k}$

In this section, we will consider the enumerations of preference sets in the set $\mathcal{P}_{n;\leq s;k}$.

Lemma 2.1. *Let $1\leq s\leq n$ and $1\leq k\leq s-1$, then*

$$p_{n;\leq s;k}=\sum_{i=1}^{s-k}\binom{n}{n-s+i+k}p_{s-i-k,s-i-1;\leq s-i-1}p_{n-s+i+k;\leq i}$$

Proof. For any $\alpha=(a_1,\ldots,a_n)\in\mathcal{P}_{n;\leq s;k}$, suppose that the $(n-i)$-th parking space is the last empty one. Obviously, $n-s+1\leq i\leq n-k$. Let $S=\{j\mid a_j>n-i\}$ and α_S be a subsequence of α determined by the subscripts in S. Then $|S|=k+i$. Let $T=[n]\setminus S$, then $|T| = n-k-i$ and $a_j < n-i$ for any $j \in T$. Let α_T be a subsequence of α determined by the subscripts in T, then $\alpha_T \in \mathcal{P}_{n-k-i,n-i-1;\leq n-i-1}$. Suppose $\alpha_s = (b_1, \ldots, b_{k+i})$, then $(b_1 - n + i, \ldots, b_{k+i} - n + i) \in \mathcal{P}_{i+k;\leq s+i-n}$.

There are $\binom{n}{i+k}$ ways to choose $i+k$ numbers from $[n]$ for the elements in S. There are $p_{i+k;\leq s+i-n}$ and $p_{n-k-i,n-i-1;\leq n-i-1}$ possibilities for α_S and α_T, respectively. Hence, for any $1 \leq k \leq s-1$, we have,

$$\begin{aligned} p_{n;\leq s;k} &= \sum_{i=n-s+1}^{n-k} \binom{n}{i+k} p_{n-i-k,n-i-1;\leq n-i-1} p_{i+k;\leq s+i-n} \\ &= \sum_{i=1}^{s-k} \binom{n}{n-s+i+k} p_{s-i-k,s-i-1;\leq s-i-1} p_{n-s+i+k;\leq i}. \end{aligned}$$

■

3 Triple-labelled rooted forests and a bijection

In this section, we consider the triplet-labelled rooted forests and bijections associated with the parking functions in the set $\mathcal{P}_{n,n+k;\leq n+k}$. Using these bijections, we give the enumerations of parking functions in the set $\mathcal{P}^{l}_{n,n+k;\leq n+k}$.

Let $\mathcal{B}_{n,k}$ be a set of all sequences $(T_0, \ldots, T_k)$ of length $k+1$ such that (1) the union of the vertex sets of $T_0, \ldots, T_k$ is $\{R_i \mid 0 \leq i \leq k\} \cup [n]$, where $R_i \notin [n]$ is just an artificial label; (2) each T_i is a tree with root R_i; (3) T_i and T_j are disjoint if $i \neq j$.

Let $F \in \mathcal{B}_{n,k}$. For any $x \in [n]$, there is an unique root R_i which is connected with x. Define the height of x to be the number of edges connecting x with root R_i. If the height of a vertex z is less than the height of x and $\{z, x\}$ is an edge of F, then z is the predecessor of x, x is a child of z, and write $z = \text{pre}(x)$ and $x \in \text{child}(z)$.

Fixing a sequence F of rooted trees in $\mathcal{B}_{n,k}$, we define a linear order $<_F$ on $[n]$ by the following rules. Let $x, y \in [n]$.

(1) For any $i \neq j$, $x \in T_i$ and $y \in T_j$, if $i < j$, then $x <_F y$.

(2) For any i and $x, y \in T_i$, if the height of x is less than the height of y, then $x <_F y$; if the height of x is equals the height of y, and $\text{pre}(x) <_F \text{pre}(y)$, then $x <_F y$.

(3) For any i and $x, y \in T_i$, if $\text{pre}(x) = \text{pre}(y)$, but $x < y$, then $x <_F y$.

The sequence formed by writing $\{1,\ldots,n\}$ in the increasing order with respect to $<_F$ is denoted by $\sigma_F^{-1}=(\sigma_F^{-1}(1),\ldots,\sigma_F^{-1}(n))$. And the permutation σ_F is the inversion of σ_F^{-1}.

Next, we define the *forest specification* of F. Let m_i be the number of the vertices in $\bigcup\limits_{j=0}^{i-1} T_j$ for $1\le i\le k+1$. Clearly, $m_{k+1}=n+k+1$. Set $\mathbf{r}_F=(r_1,\ldots,r_{n+k})$ as follows.

(1) r_1 is the number of children of the vertex R_0.

(2) r_i is the number of children of the vertex $\sigma_F^{-1}(i-1)$ if $2\le i\le m_1$.

(3) r_{m_i+1} is the number of children of the vertex R_i for $1\le i\le k$.

(4) r_i is the number of children of the vertex $\sigma_F^{-1}(i-j)$ if $m_{j-1}+2\le i\le m_j$ for some j.

Given $F\in\mathcal{B}_{n,k}$, we may obtain $(\mathbf{r}_F,\sigma_F)$. Let

$$a_i=\begin{cases}1 & \text{if } \ 1\le\sigma_F(i)\le r_1\\ k+1 & \text{if } \ 1+\sum\limits_{i=1}^{k} r_i\le\sigma_F(i)\le\sum\limits_{i=1}^{k+1} r_i\end{cases}$$

It is well known that each labeled tree on $n+1$ vertices would correspond to a parking function of length n. So, we obtain a parking function $\alpha_F=(a_1,\ldots,a_n)\in\mathcal{P}_{n,n+k;\le n+k}$ and parking spaces $m_1,\ldots,m_k$ could not be occupied.

Conversely, for any $\alpha=(a_1,\ldots,a_n)\in\mathcal{P}_{n,n+k;n+k}$, let $\mathbf{r}_\alpha=(r_1,\ldots,r_{n+k})$ be the *specificatioin* of α, i.e., $r_i=|\{j\mid a_j=i\}|$. Furthermore, we may suppose that parking spaces $m_1,\ldots,m_k$ are empty, then $r_{m_i}=0$ for $i=1,\ldots,k$.

For $1\le i\le n$, define

$$\pi_\alpha(i)=|\{a_j\mid \text{ either } a_j<a_i, \text{ or } a_j=a_i \text{ and } j<i\}|.$$

Note that $\pi_\alpha=(\pi_\alpha(1),\ldots,\pi_\alpha(n))$ is a permutation of $[n]$. Let π_α^{-1} be the inversion of π_α.

Let

$$t_i=\begin{cases}m_1-1 & \text{if } \ i=0\\ m_{i+1}-m_i-1 & \text{if } \ i=1,\ldots,k-1\\ n+k-m_k & \text{if } \ i=k\end{cases}$$

By $m_1,\ldots,m_k$, we can decompose $\mathbf{r}_\alpha$ and π_α^{-1} into the following $k+1$ subsequences,respectively:

$$\mathbf{r_i}=(r_{i,1},\ldots,r_{i,t_i})=\begin{cases}(r_1,\ldots,r_{m_1-1}) & \text{if } \ i=0\\ (r_{m_i+1},\ldots,r_{m_{i+1}-1)} & \text{if } \ i=1,\ldots,k-1\\ (r_{m_k+1},\ldots,r_{n+k)} & \text{if } \ i=k+1\end{cases}$$

and

$$\sigma_i = (\sigma_i(1), \ldots, \sigma_i(t_i)) = \begin{cases} (\pi^{-1}(1), \ldots, \pi^{-1}(m_1 - 1)) & \text{if} \quad i = 0 \\ (\pi^{-1}(m_i - i + 1), \ldots, \pi^{-1}(m_{i+1} - i - 1)) & \text{if} \quad i = 1, \ldots, k-1 \\ (\pi^{-1}(m_k - k + 1), \ldots, \pi^{-1}(n)) & \text{if} \quad i = k+1 \end{cases}$$

We associate $(\mathbf{r_i}, \sigma_i)$ with a rooted tree T_i on $t_i + 1$ vertices. The vertex set of T_i is $\{R_i\} \cup \{\sigma_i(j) \mid 1 \leq j \leq t_i\}$, where $R_i \notin [n]$ is just an artificial label. Let R_i be the root of T_i. The children of R_i are $\sigma_i(1), \ldots, \sigma_i(r_{i,1})$. For any $1 \leq j \leq t_i$, the children of $\sigma_i(j)$ are $\sigma_i(1 + \sum\limits_{m=0}^{j} r_{i,m}), \ldots, \sigma_i(\sum\limits_{m=0}^{j+1} r_{i,m})$. So, we obtain a rooted forest F_α with $k+1$ components $T_0, \ldots, T_k$ and the vertex set of F_α is $\{R_i \mid 0 \leq i \leq k\} \cup [n]$.

Lemma 3.1. *There is a bijection* ϕ *between* $\mathcal{P}_{n,n+k;\leq n+k}$ *and* $\mathcal{B}_{n,k}$.

Corollary 3.1. *Let* $\alpha = (a_1, \ldots, a_n) \in \mathcal{P}_{n,n+k;\leq n+k}$ *and* $F = (T_0, \ldots, T_k) = \phi(\alpha) \in \mathcal{B}_{n,k}$. *Suppose* $x \in [n]$, x *is a child of root* R_s *for some* $0 \leq s \leq k$. *Let* μ *be the number of the non-root vertices of* $\bigcup\limits_{j=0}^{s-1} T_j$, *then* $a_x = \mu + s + 1$.

Proof. Since $\alpha \in \mathcal{P}_{n,n+k;\leq n+k}$, we could obtain $(\mathbf{r}_\alpha, \pi_\alpha^{-1})$ and suppose that parking spaces m_1,. . .,m_k couldn't be occupied. Observe that $a_j = m_i + 1$ for some $j \in [n]$ if and only if j is a child of R_i in F. Hence, we have $a_x = m_s + 1$. So $a_x = \mu + s + 1$ since $\mu = m_s - s$. ■

Corollary 3.2. *Let* $\alpha \in \mathcal{P}_{n,n+k;\leq n+k}$ *and* $F = (T_0, \ldots, T_k) = \phi(\alpha) \in \mathcal{B}_{n,k}$. *Suppose* $x_1, x_2 \in [n]$, x_1x_2 *is an edge of* T_s *for some* $0 \leq s \leq k$ *and* $\pi_\alpha(x_1) < \pi_\alpha(x_2)$, *then* $a_{x_2} = \pi_\alpha(x_1) + s + 1$.

Proof. Since $\alpha \in \mathcal{P}_{n,n+k;\leq n+k}$, we could obtain $(\mathbf{r}_\alpha, \pi_\alpha^{-1})$ and suppose that parking spaces m_1,. . .,m_k couldn't be occupied. Then there are the subsequences $(\mathbf{r_s}, \sigma_s)$ of $(\mathbf{r}_\alpha, \pi_\alpha^{-1})$ such that $x_1 = \sigma_s(j)$ and $x_2 = \sigma_s(l + \sum\limits_{m=0}^{j} r_{i,m})$ for some j and l. Note that $\pi_\alpha(x_1) = m_s - s + j$ and $a_{x_2} = m_s + j + 1$. Hence, $a_{x_2} = \pi_\alpha(x_1) + s + 1$. ■

By Corollary 3.1 and Corollary 3.2, we may associate α with a $(k+1)$-component rooted forest $\hat{F}_\alpha$ on $n+k+1$ vertices, called *triplet-labeled rooted forest*. Since $\alpha \in \mathcal{P}_{n,n+k;\leq n+k}$, we suppose that parking spaces $m_1, \ldots, m_k$ couldn't be occupied. Let $m_0 = 0$ and $\mu_i = m_i - i$ for any $0 \leq i \leq k$. The vertex set of $\hat{F}_\alpha$ is $\{(R_i, 0, \mu_i) \mid 0 \leq i \leq k\} \cup \{(i, a_i, \pi_\alpha(i)) \mid 1 \leq i \leq n\}$

of *triplets*, where $R_i \notin [n]$ is just an artificial label for discriminating the additional triplets. Let $(R_0, 0, \mu_0), \ldots, (R_k, 0, \mu_k)$ be the root of distinct trees of $\hat{F}_\alpha$. For any two vertices $u = (x_1, y_1, z_1)$ and $v = (x_2, y_2, z_2)$, v is a child of u if there exists i such that $\mu_i \leq z_1, z_2 \leq \mu_{i+1}$ and $y_2 = z_1 + i + 1$.

For example, take $n = 12$ and $\alpha = (7, 1, 11, 12, 6, 11, 12, 2, 6, 2, 6, 12, 1) \in \mathcal{P}_{12,14;\leq 14}$. It is easy to check that $m_1 = 4$ and $m_2 = 11$. Hence, $\mu_0 = 0$, $\mu_1 = 3$ and $\mu_2 = 9$. We can obtain the following table:

i	1	2	3	4	5	6	7	8	9	10	11	12
a_i	6	1	8	12	7	12	5	8	12	2	1	5
$\pi_\alpha(i)$	6	1	8	10	7	11	4	9	12	3	2	5
$\pi_\alpha^{-1}(i)$	2	11	10	7	12	1	5	3	8	4	6	9

and $\mathbf{r}_\alpha = (2, 1, 0, 0, 2, 1, 1, 2, 0, 0, 0, 3, 0, 0)$. The rooted forest F_α associated with α is shown on Fig.1.



Fig.1. A rooted forest F_α associated with the parking function

$\alpha = (7, 1, 11, 12, 6, 11, 12, 2, 6, 2, 6, 12, 1)$.

The triplet-labelled rooted forest $\hat{F}_\alpha$ associated with α is shown on Fig.2.



Fig.2. A triplet-labelled rooted forest $\hat{F}_\alpha$ associated with the parking function

$\alpha = (7, 1, 11, 12, 6, 11, 12, 2, 6, 2, 6, 12, 1)$.

For any $\alpha \in \mathcal{P}_{n,n+k;\leq n+k}$, by triplet-labelled rooted forest $\hat{F}_\alpha$ associated with α, we could obtain the bijection ϕ between $\mathcal{P}_{n,n+k;\leq n+k}$ and $\mathcal{B}_{n,k}$.

Given $\alpha \in \mathcal{P}_{n,n+k;\leq n+k}$, let $\hat{F}_\alpha$ be a triplet-labelled rooted forest $\hat{F}_\alpha$ associated with α. Let $\psi(x,y,z) = x$ for any $(x,y,z) \in \{(R_i, 0, \mu_i) \mid 0 \leq i \leq k\} \cup \{(i, a_i, \pi_\alpha(i)) \mid 1 \leq i \leq n\}$, and two vertices $\psi(x_1,y_1,z_1)$ and $\psi(x_2,y_2,z_2)$ are adjacent if and only if (x_1,y_1,z_1) and (x_2,y_2,z_2) are adjacent in $\hat{F}$. So, we get a labeled forest F_α. Let $\phi(\alpha) = F_\alpha$, clearly, $\phi(\alpha) \in \mathcal{B}_{n,k}$.

To describe ϕ^{-1}, for each $F \in \mathcal{B}_{n,k}$, let $F = (T_0, \ldots, T_k)$ and the tree T_i have the root R_i for each i. We express F in a form, called *canonical form*, of a plane rooted forest. Let T_0, …, T_k be placed from left to right. If a vertex has more than one child then the labels of these children are increasing from left to right. Let $\mu_0 = 0$ and μ_i be the number of the non-root vertices in $\bigcup_{j=0}^{i-1} T_j$ for $1 \leq i \leq k$. Let $\theta(R_i) = (R_i, 0, \mu_i)$ for each i. For any $j \in [n]$, let $\theta(j) = (j, y_j, z_j)$, where y_j and z_j are determined by the following algorithm A.

Algorithm A.

(1) Let $F = (T_0, \ldots, T_k)$. Traverse T_i by a breadth-first search, suppose j is the s-th non-root vertex and label the third entries $z_j = s + \mu_i$.

(2) For any two $v = (x_1, y_1, z_1)$ and $u = (x_2, y_2, z_2)$, if u is a child of v and $x_1 = R_i$, then $y_2 = \mu_i + i + 1$; if u is a child of v, the vertices x_1 and x_2 are in T_i and $x_1, x_2 \in [n]$, then $y_2 = z_1 + i + 1$.

Note that if u is a child v in T_i, then $y_2 - \mu_i - i = z_2 - \mu_i = 1$ if $x_1 = R_i$; and $z_2 > z_1$ implies that $y_2 - \mu_i - i = (z_1 + i + 1) - \mu_i - i \leq z_2 - \mu_i$ if $x_1 \neq R_i$. Sorting the triplets of non-vertices by the first entries, the sequence $\varphi^{-1}(F) = (y_1, \ldots, y_n)$, which is formed by their second entries, is the required parking function. Obviously, parking spaces $\mu_i + i$ couldn't be occupied for $1 \leq i \leq k$. ■

For any $\alpha = (a_1, \ldots, a_n) \in \mathcal{P}_{n,n+k;\leq n+k}$, we may obtain $\mathbf{r}_\alpha = (r_1, \ldots, r_{n+k})$ and $\pi_\alpha = (\pi_\alpha(1), \ldots, \pi_\alpha(n))$ and suppose that parking spaces $m_1, \ldots, m_k$ couldn't be occupied with $m_1 < m_2 < \ldots < m_k$, then define the following parameters about α:

(1) $l_\alpha :$ the leading term of α, i.e., $l_\alpha = a_1$;

(2) $n_\alpha :$ the number of a_1 in α, i.e., $t_\alpha = |\{i \mid a_i = a_1\}|$;

(3) $m_\alpha = m_k$;

(4) $g_\alpha = max\{m_i \mid m_i < l_\alpha\}$;

(5) $h_\alpha = max\{i \mid m_i < l_\alpha\}$;

(6) $\tau_\alpha = |\{i \mid a_j < l_\alpha\}| + 1$;

Clearly, $\tau_\alpha \geq l_\alpha - h_\alpha$. Now, let $\mathcal{P}^l_{n,n+k;\leq n+k} = \{\alpha \in \mathcal{P}_{n,n+k;\leq n+k} \mid l_\alpha = l\}$. Let $\mathcal{F}^l_{n,k}$ denote the set of triplet-labelled rooted forests F'_α associated with $\alpha \in \mathcal{P}^l_{n,n+k;\leq n+k}$. If $\mathcal{R}$ is a set consisting of some parking functions, then we always use $\mathcal{F}(\mathcal{R})$ to denote the set of triplet-labelled rooted forests associated with the parking functions in $\mathcal{R}$.

The following lemma has the same proof as the lemma in [3]. For the sake of completeness, we still prove it as follows.

Lemma 3.2. *Let $k \geq 0$. For any $1 \leq l \leq n+k-1$, let $\mathcal{A}_1 = \{\alpha \in \mathcal{P}^l_{n,n+k;\leq n+k} \mid \tau_\alpha > l - h_\alpha$ or $n_\alpha \geq 2\}$ and $\mathcal{C}_1 = \{\alpha \in \mathcal{P}^{l+1}_{n,n+k;\leq n+k} \mid l \geq g_\alpha + 1\}$. Then there is a bijection from $\mathcal{A}_1$ to $\mathcal{C}_1$.*

Proof. It suffices to establish a bijection $\psi : \mathcal{F}(\mathcal{A}_1) \to \mathcal{F}(\mathcal{C}_1)$. Given an $\hat{F}_\alpha \in \mathcal{F}(\mathcal{A}_1)$, let $u = (1, l, \pi_\alpha(1)) \in \hat{F}_\alpha$. Obviously, $u \in T_{h_\alpha}$ and $l \geq g_\alpha + 1$. If $\tau_\alpha > l - h_\alpha$, then $\pi_\alpha(1) > l - h_\alpha$ and $T_{h_\alpha} - T_{h_\alpha}(u)$ has at least $l - h_\alpha$ vertices. On the other hand, $u \in T_{h_\alpha}$ implies that there are at least $l - h_\alpha - 1$ terms a_j satisfying $g_\alpha + 1 \leq a_j < l$. Hence, if $n_\alpha \geq 2$, then there are at least $l - h_\alpha$ vertices in $T_{h_\alpha} - T_{h_\alpha}(u)$ as well.

Let $(R_{h_\alpha}, 0, g_\alpha - h_\alpha)$ be the 0-th vertex. Traverse $T_{h_\alpha} - T_{h_\alpha}(u)$ by breadth-first search and locate the $(l - g_\alpha)$-th vertex in $T_{h_\alpha} - T_{h_\alpha}(u)$, say v. By attaching $T_{h_\alpha}(u)$ to v so that u is the first child of v in $T_{h_\alpha} - T_{h_\alpha}(u)$, updating the second and the third entries of all non-root vertices by Algorithm A, and the other trees remain unchangeable, we obtain $\psi(\hat{F}_\alpha)$. the triplet of u becomes $(1, l+1, \pi(1))$. $l \geq g_\alpha + 1$ implies that $\psi(\hat{F}_\alpha) \in \mathcal{F}(\mathcal{C}_1)$.

To find ψ^{-1}, given an $\hat{F}_\beta \in \mathcal{F}(\mathcal{C}_1)$, let $u = (1, l+1, \pi_\beta(1)) \in \hat{F}_\beta$ and v the parent of u. Clearly, $u \in T_{h_\beta}$. Since $l \geq g_\beta + 1$, v isn't the root $(R_{h_\beta}, 0, g_\beta - h_\beta)$ of T_{h_β}. In $\hat{F}_\beta$, we locate the vertex, say w, the third entry of which is equal to $l - h_\beta - 1$. Attach $T_{h_\beta}(u)$ to w so that u is the first child of w. By Algorithm A, the updated triplet of u becomes $(1, l, \pi(1))$ and the other trees are unchangeable. We observe that either $\pi(1) = l$ if v is another child of w, or $\pi(1) > l$ otherwise. Hence $\psi^{-1}(\hat{F}_\beta) \in \mathcal{F}(\mathcal{R})$. ■

For example, take $n = 12$, $k = 2$ and $l = 2$. We consider a parking function $\alpha = (2, 4, 8, 12, 9, 1, 12, 8, 4, 12, 1, 3) \in \mathcal{P}_{12,14;\leq 14}$. Then $h_\alpha = 0$, $n_\alpha = 1$, $g_\alpha = 0$ and $\tau_\alpha = 3$. Observe that $\tau_\alpha > l - h_\alpha$. On Fig.3 is the forest $\hat{F}_\alpha$ associated with α. Let $u = (1, 2, 3)$. Note that $v = (11, 1, 2)$ is the 2-th vertex of $\hat{F}_\alpha - T_1(u)$ that is visited by a breadth-first search. On Fig.4 is the corresponding forest $\psi(\hat{F}_\alpha)$, which is obtained from $\hat{F}_\alpha - T_1(u)$ with $T_1(u)$ attached to v and with the second and third entries of the triplet updated. Sorting the triplets of non-root vertices by the first entries, we retrieve the corresponding parking function $\alpha = (3, 4, 8, 12, 9, 1, 12, 8, 4, 12, 1, 3)$ with leading term 3 from their second entries.



Fig.3. The forest F'_αassociated with the parking function $\alpha = (2, 4, 8, 12, 9, 1, 12, 8, 4, 12, 1, 3)$



Fig.4. The corresponding forest $\phi(F'_\alpha)$

Lemma 3.3. *Let k be an integer with $k \geq 1$. For any $1 \leq l \leq n + k - 1$, Let $\mathcal{A}_2 = \{\alpha \in \mathcal{P}^l_{n,n+k;\leq n+k} \mid \tau_\alpha = l - h_\alpha, n_\alpha = 1, h_\alpha \leq k - 1\}$ and $\mathcal{C}_2 = \{\alpha \in \mathcal{P}^{l+1}_{n,n+k;\leq n+k} \mid l = g_\alpha\}$, then there is a bijection from $\mathcal{A}_2$ to $\mathcal{C}_2$.*

Proof. It suffices to establish a bijection $\psi : \mathcal{F}(\mathcal{A}_2) \to \mathcal{F}(\mathcal{C}_2)$. Given an $\hat{F}_\alpha \in \mathcal{F}(\mathcal{A}_2)$, let $u = (1, l, \pi_\alpha(1)) \in \hat{F}_\alpha$. Obviously, $u \in T_{h_\alpha}$. Let $r_{h_\alpha+1}$ be the root of the tree $T_{h_\alpha+1}$. Delete $T_{h_\alpha}(u)$ from T_{h_α} and attach $T_{h_\alpha}(u)$ to $r_{h_\alpha+1}$ so that u is the first child of $r_{h_\alpha+1}$. Then updating the second and the third entries of all non-root vertices by Algorithm A, and the other trees remain unchangeable, we obtain $\psi(\hat{F}_\alpha)$. The triplet of u becomes $(1, l + 1, l - h_\alpha)$. Suppose $\hat{F}_\beta = \psi(\hat{F}_\alpha)$, then $g_\beta = l$, hence $\psi(\hat{F}_\alpha) \in \mathcal{F}(\mathcal{C}_2)$.

To find ψ^{-1}, given an $\hat{F}_\beta \in \mathcal{F}(\mathcal{C}_2)$, let $u = (1, l+1, \pi_\beta(1)) \in \hat{F}_\beta$. Since $l = g_\beta$, we have $h_\beta \geq 1$ and that the parent of u is the root R_{h_β} of the tree T_{h_β} in $\hat{F}_\beta$. Traverse $T_{h_\beta - 1}$ by breadth-first search and suppose v is the last vertex. Delete $T_{h_\beta}(u)$ from T_{h_β} and attach $T_{h_\beta}(u)$ to v. Then updating the second and the third entries of all non-root vertices by Algorithm A, and the other trees remain unchangeable, we obtain $\psi^{-1}(\hat{F}_\beta)$. The triplet of u becomes $(1, l, l - h_\beta + 1)$. Hence, $\psi^{-1}(\hat{F}_\beta) \in \mathcal{F}(\mathcal{A}_2)$. ■

For example, take $n = 10$, $k = 2$ and $l = 3$. We consider a parking function $\alpha = (3, 10, 4, 10, 7, 1, 4, 1, 10, 7) \in \mathcal{P}_{10,12;\leq 12}$. Then $h_\alpha = 0$, $\tau_\alpha = 3$ and $t_\alpha = 1$. Observe that $\tau_\alpha = l - h_\alpha$. On Fig.6 is the corresponding forest $\psi(\hat{F}_\alpha)$, which is obtained from $\hat{F}_\alpha - T_1(u)$ with $T_1(u)$ attached to $(R_1, 0, 5)$ and with the second and third entries of the triplet updated. Sorting the triplets of non-root vertices by the first entries, we retrieve the corresponding parking function $(4, 10, 5, 10, 4, 1, 5, 1, 10, 4)$ with leading term 4 from their second entries.



Fig.5. The forest F'_αassociated with the parking function $\alpha = (3, 10, 4, 10, 7, 1, 4, 1, 10, 7)$



Fig.6. The corresponding forest $\phi(F'_\alpha)$

Lemma 3.4. *Let $k \geq 0$. For any $k+1 \leq l \leq n+k$, Let $\mathcal{A}_3 = \{\alpha \in \mathcal{P}^l_{n,n+k;\leq n+k} \mid \tau_\alpha = l - h_\alpha, n_\alpha = 1, h_\alpha = k\}$ and $\mathcal{C}_3 = \{\beta \in \mathcal{P}^{n+k+1}_{n,n+k+1;\leq n+k+1} \mid m_\beta = l\}$, then there is a bijection from $\mathcal{A}_3$ to $\mathcal{C}_3$.*

Proof. For any $\alpha = (l, a_2, \ldots, a_n) \in \mathcal{A}_3$, let $\beta = (n+k+1, a_2, \ldots, a_n)$. Let $S = \{j \mid l+1 \leq a_i \leq n+k\}$ and $T = \{j \mid m_\alpha + 1 \leq a_i \leq l-1\}$. Furthermore, let α_S and α_T be two subsequences of α determined by the subscripts in S and T, respectively. Since $\alpha \in \mathcal{A}_3$, α_S

and α_T correspond with a parking function of length $n+k-l$ and $l-m_\alpha-1$, respectively. So, $h_\alpha = k$ implies $\beta \in \mathcal{C}_3$.

Conversely, for any $\beta = (n+k+1, a_2, \ldots, a_n) \in \mathcal{C}_3$, let $\alpha = (l, a_2, \ldots, a_n)$. $m_\beta = l$ implies $\alpha \in \mathcal{A}_3$. This complete the proof. ■

Lemma 3.5. *Let $k \geq 0$. For any $k+1 \leq l \leq n+k$, Let $\mathcal{A}_3 = \{\alpha \in \mathcal{P}^l_{n,n+k;\leq n+k} \mid \tau_\alpha = l - h_\alpha, n_\alpha = 1, h_\alpha = k\}$, then $|\mathcal{A}_3| = \binom{n-1}{l-k-1} p_{l-k-1,l-1,\leq l-1} p_{n+k-l}$.*

Proof. Lemma 3.4 implies that $|\mathcal{A}_3| = |\mathcal{C}_3|$. For any $\alpha = (a_1, \ldots, a_n) \in \mathcal{C}_3$, let $S = \{j \mid a_i \leq l-1\}$ and $T = [n] \setminus S$, then $|S| = l-k-1$ and $|T| = n+k+1-l$. Furthermore, let α_S and α_T be two subsequences of α determined by the subscripts in S and T, respectively. Obviously, $\alpha_S \in \mathcal{P}_{l-k-1,l-1,\leq l-1}$. Suppose $\alpha_T = (b_1, \ldots, b_{n+k+1-l})$, since $l+1 \leq b_j \leq n+k+1$ for any $1 \leq j \leq n+k+1-l$, we have $(b_1 - l, \ldots, b_{n+k+1-l} - l) \in \mathcal{P}^{n+k+1-l}_{n+k+1-l}$.

There are $\binom{n-1}{l-k-1}$ ways to choose $l-k-1$ numbers from $[2,n]$ for the elements in S. There are $p_{l-k-1,l-1,\leq l-1}$ and p_{n+k-l} possibilities for α_S and α_T, respectively. Hence, we have $|\mathcal{A}_3| = |\mathcal{C}_3| = \binom{n-1}{l-k-1} p_{l-k-1,l-1,\leq l-1} p_{n+k-l}$. ■

4 Enumerations for parking functions in $\mathcal{P}^l_{n,n+k;\leq n+k}$

In this section, with the benefit of the triplet-labelled rooted forests, we enumerate parking functions in the set $\mathcal{P}^l_{n,n+k;\leq n+k}$.

Theorem 4.1. *For any $k \geq 0$, we have $p^1_{n,n+k;\leq n+k} = p^2_{n,n+k;\leq n+k} = \ldots = p^{k+1}_{n,n+k;\leq n+k}$.*

Proof. Let $\mathcal{A}_1$ and $\mathcal{C}_1$ be defined as that in Lemma 3.2, $\mathcal{A}_2$ and $\mathcal{C}_2$ defined as that in Lemma 3.3. Note that $\mathcal{P}^l_{n,n+k;\leq n+k} = \bigcup_{i=1}^{2} \mathcal{A}_i$ and $\mathcal{P}^{l+1}_{n,n+k;\leq n+k} = \bigcup_{i=1}^{2} \mathcal{C}_i$. Hence, by Lemmas 3.2 and 3.3, we obtain the desired results. ■

Theorem 4.2. *(1) $p^1_{n,n+k;\leq n+k} = p_{n-1,n+k;\leq n+k}$ for any $k \geq 0$ and $n \geq 1$;*

(2) $p^1_{n,n+k,\leq n+k} = \sum_{i=1}^{n} \binom{n-1}{i-1} p^1_{i,i+k-1,\leq i+k-1} p_{n-i}$ for any $k \geq 1$.

Proof. (1) For any $\alpha = (1, a_2, \ldots, a_n) \in \mathcal{P}^1_{n,n+k;\leq n+k}$, let $\psi(\alpha) = (a_2, \ldots, a_n)$, then $\psi(\alpha) \in \mathcal{P}_{n-1,n+k;\leq n+k}$. Obviously, the mapping ψ is a bijection between the sets $\mathcal{P}^1_{n,n+k;\leq n+k}$ and $\mathcal{P}_{n-1,n+k;\leq n+k}$. Hence, $p^1_{n,n+k;\leq n+k} = p_{n-1,n+k;\leq n+k}$ for any $k \geq 0$ and $n \geq 1$.

(2) For any $\alpha \in \mathcal{P}^1_{n,n+k,\leq n+k}$, we suppose that the last parking space which don't be occupied is i. Obviously, $k+1 \leq i \leq n+k$. Let $S = \{j \mid a_j \leq i-1, a_j \in \alpha\}$ and $T = [n] \setminus S$, then $|S| = i-k$, $|T| = n+k-i$. Let α_S and α_T be two subsequences of α determined by the subscripts in S and T, respectively. Then we have $\alpha_S \in \mathcal{P}^1_{i-k,i-1,\leq i-1}$. Suppose $\alpha_T = (b_1, \ldots, b_{n+k-i})$, then $(b_1 - i, \ldots, b_{n+k-i} - i) \in \mathcal{P}_{n+k-i}$.

There are $\binom{n-1}{i-k-1}$ ways to choose $i-k-1$ numbers from $[2,n]$ for elements in S since $1 \in S$. There are $p^1_{i-k,i-1,\leq i-1}$ and p_{n+k-i} possibilities for the preference sets α_S and α_T, respectively.

Hence, we have

$$\begin{aligned} p^1_{n,n+k,\leq n+k} &= \sum_{i=k+1}^{n+k} \binom{n-1}{i-k-1} p^1_{i-k,i-1,\leq i-1} p_{n+k-i} \\ &= \sum_{i=1}^{n} \binom{n-1}{i-1} p^1_{i,i+k-1,\leq i+k-1} p_{n-i}. \end{aligned}$$

■

Theorem 4.3. *Let $n \geq 1$, $k \geq 0$ and $k+1 \leq l \leq n+k-1$. Then*

$$p^l_{n,n+k;\leq n+k} - p^{l+1}_{n,n+k;\leq n+k} = \binom{n-1}{l-k-1} p_{l-k-1,l-1;\leq l-1} p_{n+k-l}.$$

Proof. Let $\mathcal{A} = \{\alpha \in \mathcal{P}^l_{n,n+k;\leq n+k} \mid h_\alpha = k, n_\alpha = 1, \tau_\alpha = l-k\}$ and $\mathcal{B} = \mathcal{P}^l_{n,n+k;\leq n+k} \setminus \mathcal{A}$. By Lemmas 3.2 and 3.3, there is a bijection from the sets $\mathcal{B}$ to $\mathcal{P}^{l+1}_{n,n+k;\leq n+k}$. Hence, $p^l_{n,n+k;\leq n+k} - p^{l+1}_{n,n+k;\leq n+k} = |\mathcal{A}|$. Lemma 3.5 implies that

$$p^l_{n,n+k;\leq n+k} - p^{l+1}_{n,n+k;\leq n+k} = \binom{n-1}{l-k-1} p_{l-k-1,l-1;\leq l-1} p_{n+k-l}.$$

■

Lemma 4.1. *For any $k \geq 0$ and $n \geq 1$, we have $p^{n+k}_{n,n+k;\leq n+k} = p_{n-1,n+k-1;\leq n+k-1}$*

Proof. For any $\alpha = (n+k, a_2, \ldots, a_n) \in \mathcal{P}^{n+k}_{n,n+k;\leq n+k}$, let $\varphi(\alpha) = (a_2, \ldots, a_n)$, then $\varphi(\alpha) \in \mathcal{P}_{n-1,n+k-1;\leq n+k-1}$. Obviously, this is a bijection. ■

5 The Bijections for k-flaw preference sets in $\mathcal{P}^l_{n;\leq s;k}$

In this section, we consider triplet-labelled rooted forests associated with preference sets in the set $\mathcal{P}^l_{n;\leq s;k}$.

For any $1 \leq s \leq n$, let $\mathcal{B}_{n,s,k}$ denote a set of the forests $F = (T_0, \ldots, T_k) \in \mathcal{B}_{n,k}$ such that (1) the tree T_k has at least $n + k - s + 1$ vertices; (2) all the vertices $\sigma_F^{-1}(s - k), \sigma_F^{-1}(s + 1 - k), \ldots, \sigma_F^{-1}(n)$ are leaves, where σ_F^{-1} is the linear order on $[n]$ with respect to $<_F$.

Lemma 5.1. *Let n and k be two nonnegative integers. Suppose that $1 \leq s \leq n$ and the mapping ϕ is defined as that in Lemma 3.1, then ϕ is a bijection between the sets $\mathcal{P}_{n;\leq s;k}$ and $\mathcal{B}_{n,s,k}$.*

Proof. For any $\alpha = (a_1, \ldots, a_n) \in \mathcal{P}_{n;\leq s;k}$, if we add k parking spaces, then α can be viewed as an element in the set $\mathcal{P}_{n,n+k;\leq n+k}$. So, we may suppose that the empty parking spaces are $m_1, \ldots, m_k$ and $\hat{F}_\alpha = (T_0, \ldots, T_k)$ is the triple-let labeled rooted forest associated with α. Lemma 3.1 implies that $\phi(\alpha) \in \mathcal{B}_{n,k}$. Since $s \leq n$, we have $m_k \leq s-1$, hence, the tree T_k has at least $n + k - s + 1$ vertices. On the other hand, let $\mathbf{r}_\alpha$ and π_α be the specification and the permutation on $[n]$ of α, respectively. Then the vertex $\pi_\alpha^{-1}(i)$ is a leaf in T_k since $r_{i+k+1} = 0$ for any $i \in [s - k, n]$. So, $\phi(\alpha) \in \mathcal{B}_{n,s,k}$.

Conversely, for any $F \in \mathcal{B}_{n,s,k}$, Lemma 3.1 tells us that $\phi^{-1}(F) = (a_1, \ldots, a_n) \in \mathcal{P}_{n,n+k;\leq n+k}$. Suppose that all the empty parking spaces are $m_1, \ldots, m_k$, then $m_k \leq s - 1$ since T_k has at least $n + k - s + 1$ vertices. Let σ_F^{-1} be the linear order on $[n]$ with respect to $<_F$. Furthermore, let $\hat{F}$ be the triple-let labeled rooted forest from F. Then $a_i \leq s$ for any $i \in [n]$ since the vertices $\sigma_F^{-1}(s-k), \sigma_F^{-1}(s-k+1), \ldots, \sigma_F^{-1}(n)$ are leaves. If we erase exact k parking spaces $n+1, \ldots, n+k$, then there are k cars which can't be parked, hence, $\phi^{-1}(F) \in \mathcal{P}_{n;\leq s;k}$. ■

The proofs of the following four lemmas is similar to Lemmas 3.2, 3.3, 3.4 and 3.5. We just state them as follows.

Lemma 5.2. *Suppose that $k \geq 1$, $l \geq 1$ and $l + 1 \leq s \leq n$. Let $\mathcal{A}_1' = \{\alpha \in \mathcal{P}_{n;\leq s;k}^{l} \mid h_\alpha \leq k - 1$ and $\tau_\alpha > l - h_\alpha$ or $n_\alpha \geq 2\}$ and $\mathcal{C}_1' = \{\alpha \in \mathcal{P}_{n;\leq s;k}^{l+1} \mid h_\alpha \leq k - 1$ and $l \geq g_\alpha + 1\}$. Then there is a bijection from $\mathcal{A}_1'$ to $\mathcal{C}_1'$.*

Lemma 5.3. *Suppose that $k \geq 2$ and $1 \leq l \leq s - 3$. Let $\mathcal{A}_2' = \{\alpha \in \mathcal{P}_{n;\leq s;k}^{l} \mid \tau_\alpha = l - h_\alpha, n_\alpha = 1, h_\alpha \leq k - 2\}$ and $\mathcal{C}_2' = \{\alpha \in \mathcal{P}_{n;\leq s;k}^{l+1} \mid h_\alpha \leq k - 1$ and $l = g_\alpha\}$, then there is a bijection from $\mathcal{A}_2'$ to $\mathcal{C}_2'$.*

Lemma 5.4. *Suppose that* $k \geq 1$ *and* $k+1 \leq l \leq s-2$*. Let* $\mathcal{A}'_3 = \{\alpha \in \mathcal{P}^l_{n;\leq s;k} \mid \tau_\alpha = l - h_\alpha, n_\alpha = 1, h_\alpha = k-1\}$ *and* $\mathcal{C}'_3 = \{\beta \in \mathcal{P}^s_{n;\leq s;k+1} \mid m_k = l\}$*, then there is a bijection from* $\mathcal{A}'_3$ *to* $\mathcal{C}'_3$*.*

Lemma 5.5. *Suppose* $k \geq 1$ *and* $k+1 \leq l \leq s-2$*. Let* $\mathcal{A}'_3 = \{\alpha \in \mathcal{P}^l_{n;\leq s;k} \mid \tau_\alpha = l - h_\alpha, n_\alpha = 1, h_\alpha = k-1\}$*, then* $|\mathcal{A}'_3| = \binom{n-1}{l-k} p_{l-k,l-1,\leq l-1} p^{s-l}_{n+k-l;\leq s-l;1}$*.*

Using the above four lemmas, we may consider the problem of enumerations of preferences sets in the set $\mathcal{P}^l_{n;\leq s;k}$.

6 Enumerations for the preference sets in $\mathcal{P}^l_{n;\leq s;k}$

First, we study the case with $1 \leq l \leq k$.

Theorem 6.1. *Suppose that* $1 \leq s \leq n$*. For any* $k \geq 1$*, we have* $p^1_{n;\leq s;k} = p^2_{n;\leq s;k} = \ldots = p^k_{n;\leq s;k}$*.*

Proof. Let $\mathcal{A}'_1$ and $\mathcal{C}'_1$ be defined as that in Lemma 5.2, $\mathcal{A}'_2$ and $\mathcal{C}'_2$defined as that in Lemma 5.3. Note that $\mathcal{P}^l_{n;\leq s;k} = \bigcup_{i=1}^{2} \mathcal{A}'_i$ and $\mathcal{P}^{l+1}_{n;\leq s;k} = \bigcup_{i=1}^{2} \mathcal{C}'_i$. Hence, by Lemmas 5.2 and 5.3, we obtain the desired results. ■

Theorem 6.2. *Let* $k \geq 1$ *and* $k+1 \leq s \leq n$*, then*

$$p^1_{n;\leq s;k} = \sum_{i=1}^{s-k-1} \binom{n-1}{s-i-k-1} p_{n+k-s+i;\leq i} p^1_{s-k-i,s-i-1;\leq s-i-1}.$$

Proof. For any $\alpha = (a_1, \ldots, a_n) \in \mathcal{P}^1_{n;\leq s;k}$, we suppose that the last empty parking space is $s-i$. Obviously, $1 \leq i \leq s-k-1$. Let $S = \{j \mid s-i+1 \leq a_j \leq s\}$ and α_S be a subsequence of α determined by the subscripts in S, then $|S| = n+k-s+i$. Suppose $\alpha_S = (b_1, \ldots, b_{n+k-s+i})$, then $(b_1 - s + i, \ldots, b_{n+k-s+i} - s + i) \in \mathcal{P}_{n+k-s+i;\leq i}$. Let $T = [n] \setminus S$ and α_T be a subsequence of α determined by the subscripts in T, then $|T| = s-k-i$ and $\alpha_T \in \mathcal{P}^1_{s-k-i,s-i-1;\leq s-i-1}$.

There are $\binom{n-1}{s-k-i-1}$ ways to choose $s-k-i-1$ numbers from $[2,n]$ for the elements in T since $1 \in T$. There are $p_{n+k-s+i}$ and $p^1_{s-k-i,s-i-1;\leq s-i-1}$ possibilities for the parking function α_S and the preference set α_T, respectively. Hence, we have

$$p^1_{n;\leq s;k} = \sum_{i=1}^{s-k-1} \binom{n-1}{s-i-k-1} p_{n+k-s+i;\leq i} p^1_{s-k-i,s-i-1;\leq s-i-1}.$$

■

Now, we consider the case with $l = s$.

Theorem 6.3. *$p^s_{n;\leq s;k+1} = p^1_{n;\leq s;k}$ for any $k \geq 1$ and $k+2 \leq s \leq n$.*

Proof. It suffices to establish a bijection $\varphi : \mathcal{F}(\mathcal{P}^1_{n;\leq s;k}) \to \mathcal{F}(\mathcal{P}^s_{n;\leq s;k+1})$. Given an $\hat{F}_\alpha = (T_0, \ldots, T_k) \in \mathcal{F}(\mathcal{P}^1_{n,\leq s,k})$, let $u = (1, 1, \pi_\alpha(1)) \in \hat{F}_\alpha$. Clearly, $u \in T_0$. Deleting $T_0(u)$ from T_0, we denote $T'_0 = T_0 - T_0(u)$. Suppose the number of the non-root vertices in T'_0 is a, let the triplet of u becomes $(R_1, 0, a)$ and $T'_1 = T_0(u)$. Let the triplet $(R_2, 0, \mu_2)$ of the root in T_2 become $(R_3, 0, \mu_2 - 1)$ and let $T'_3 = T_2$. For any $j \geq 2$, let the triplet $(R_i, 0, \mu_i)$ of the root in T_i become $(R_{i+1}, 0, \mu_i)$ and let $T'_{i+1} = T_i$. Traverse T'_{k+1} by breadth-first search and suppose v is the $(s-1-k-\mu_k)$-th vertex, attach a new vertex w, where the first entry of w is 1, as the first child v. Then updating the second and the third entries of all non-root vertices by Algorithm A, we obtain $\varphi(\hat{F}_\alpha) \in \mathcal{F}(\mathcal{P}^s_{n;\leq s;k+1})$.

To find φ^{-1}, given an $\hat{F}_\beta = (T'_0, \ldots, T'_{k+1}) \in \mathcal{F}(\mathcal{P}^s_{n;\leq s;k+1})$, let $w = (1, s, \pi_\beta(1)) \in \hat{F}_\beta$. Clearly, $w \in T'_{k+1}$. Delete w from T'_{k+1}. Suppose that the root of T'_1 is $u = (R_1, 0, \mu_1)$. Let the first entry of the triplet of u become 1 and attach u to be the first child of the root of T'_0. We denote the obtained tree as T_0. Let the triplet $(R_2, 0, \mu_2)$ of the root in T'_2 become $(R_1, 0, \mu_2 + 1)$ and let $T_1 = T'_2$. For any $j \geq 3$, let the triplet $(R_i, 0, \mu_i)$ of the root in T'_i become $(R_{i-1}, 0, \mu_i)$ and let $T'_{i-1} = T_i$. Then updating the second and the third entries of all non-root vertices by Algorithm A, we obtain $\varphi^{-1}(\hat{F}_\alpha) \in \mathcal{F}(\mathcal{P}^1_{n;\leq s;k})$. ■

Theorem 6.4. *Let $k \geq 0$ and $k+1 \leq s \leq n$. Then $p^s_{n+1;\leq s;k} = p_{n;\leq s;k}$.*

Proof. For any $\beta = (s, b_1, \ldots, b_n) \in \mathcal{P}^s_{n+1,\leq s,k}$, we consider the mapping $\varphi(\beta) = (b_1, \ldots, b_n)$. It is easy to check that φ if a bijection from the sets $p^s_{n+1;\leq s;k}$ to $\mathcal{P}_{n;\leq s;k}$. Hence, $p^s_{n+1;\leq s;k} = p_{n;\leq s;k}$. ■

Corollary 6.1. *Let k, n and s be three integers. Suppose $k \geq 1$ and $k+2 \leq s \leq n$. Then $p^1_{n+1;\leq s;k} = p_{n;\leq s;k+1}$.*

Proof. By Theorems 6.3 and 6.4, we immediately obtain the desired results. ■

Theorem 6.5. *$p^n_{n;1} = p^2_n$ for any $n \geq 2$.*

Proof. It suffices to establish a bijection $\varphi : \mathcal{F}(\mathcal{P}_n^2) \to \mathcal{F}(\mathcal{P}_{n,1}^n)$. Given an $\hat{T}_\alpha \in \mathcal{F}(\mathcal{P}_n^2)$, let $u = (1, 2, \pi_\alpha(1)) \in \hat{T}_\alpha$ and $w = (R_0, 0, 0)$ be the root of $\hat{T}_\alpha$. Suppose the number of the vertices in $\hat{T}_\alpha(u)$ is a. Obviously, $a \geq 1$. Let the triplets of w and u become $(R_1, 0, a-1)$ and $(R_0, 0, 0)$, respectively. We denote $T_0 = \hat{T}_\alpha(u)$. Deleting $\hat{T}_\alpha(u)$ from $\hat{T}_\alpha$ and traversing $\hat{T}_\alpha - \hat{T}_\alpha(u)$ by breadth-first search and suppose v is the $(n-1-a)$-th vertex, attaching a new vertex s, where the first entry of s is 1, as the first child v, we denote the obtained tree as T_1. Then updating the second and the third entries of all non-root vertices by Algorithm A, we obtain $\psi(\hat{T}_\alpha) \in \mathcal{F}(\mathcal{P}_{n;1}^n)$.

To find φ^{-1}, given an $\hat{F}_\beta \in \mathcal{F}(\mathcal{P}_{n;1}^n)$, let $s = (1, n, \pi_\beta(1)) \in \hat{F}_\beta = (T_0, T_1)$. Clearly, $s \in T_1$. Suppose $u = (R_0, 0, 0)$ is the root of T_0, the number of non-root vertices of T_0 is a and the root of T_1 is $w = (R_1, 0, a)$. Delete the vertex s from T_1. Let the first entry of the triplets of u become 1 and the triplets of w become $(R_0, 0, 0)$. Traverse T_1 by breadth-first search and suppose v is the first non-root vertex and attach u to be the first child of v. Then updating the second and the third entries of all non-root vertices by Algorithm A, we obtain $\varphi^{-1}(\hat{F}_\beta)$. The triplet of u becomes $(1, 2, \pi(1))$. Hence, $\varphi^{-1}(\hat{F}_\beta) \in \mathcal{F}(\mathcal{P}_n^2)$. ■

Lemma 6.1. *For any $n \geq 2$, we have $p_{n;1}^n - p_{n;1}^{n-1} = p_{n-2}$*

Proof. Let $\mathcal{A}_1 = \{\alpha \in \mathcal{P}_{n;1}^{n-1} \mid r_{n-1} + r_n \leq 2\}$ and $\mathcal{C}_1 = \{\beta \in \mathcal{P}_{n;1}^n \mid r_{n-1} + r_n \leq 2 \text{ and } m_\beta < n-1\}$. Let $\mathcal{A}_2 = \{\alpha \in \mathcal{P}_{n;1}^{n-1} \mid r_{n-1} = 3\}$ and $\mathcal{C}_2 = \{\beta \in \mathcal{P}_{n;1}^n \mid r_{n-1} = 2 \text{ and } r_n = 1\}$. For any $\alpha = (n-1, a_2, \ldots, a_n) \in \mathcal{A}_1 \cup \mathcal{A}_2$, we consider the mapping $\psi_1(\alpha) = (n, a_2, \ldots, a_n) \in \mathcal{C}_1 \cup \mathcal{C}_2$. Clearly, the mapping ψ_1 is a bijection from the sets $\mathcal{A}_1 \cup \mathcal{A}_2$ and $\mathcal{C}_1 \cup \mathcal{C}_2$.

Let $\mathcal{A}_3 = \{\alpha \in \mathcal{P}_{n;1}^{n-1} \mid r_n = 1 \text{ and } r_{n-1} = 2\}$ and $\mathcal{C}_3 = \{\beta \in \mathcal{P}_{n;1}^n \mid r_{n-1} = 1 \text{ and } r_n = 2\}$. For any $\alpha = (a_1, \ldots, a_n) \in \mathcal{A}_3$, obviously, $a_1 = n-1$ and there are $i \neq 1$ and j such that $a_i = n-1$ and $a_j = n$. Let $b_1 = b_i = n$, $b_j = n-1$ and $b_h = a_h$ for any $h \notin \{1, i, j\}$, then $\beta = (b_1, \ldots, b_n) \in \mathcal{C}_3$. It is easy to obtain that ψ_2 is a bijection between the sets $\mathcal{A}_3$ to $\mathcal{C}_3$.

Let $\mathcal{A}_4 = \{\alpha \in \mathcal{P}_{n;1}^{n-1} \mid r_n = 2 \text{ and } r_{n-1} = 1\}$ and $\mathcal{C}_4 = \{\beta \in \mathcal{P}_{n;1}^n \mid r_n = 2 \text{ and } m_\beta = n-1\}$. Note that $(\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3, \mathcal{A}_4)$ and $(\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{C}_4)$ form a partition for $\mathcal{P}_{n;1}^{n-1}$ and $\mathcal{P}_{n;1}^n$, respectively. Hence, $p_{n;1}^n - p_{n;1}^{n-1} = |\mathcal{P}_{n;1}^n| - |\mathcal{P}_{n;1}^{n-1}| = |\mathcal{C}_4| - |\mathcal{A}_4|$. It is easy to obtain that $|\mathcal{C}_4| = (n-1)p_{n-2}$ and $|\mathcal{A}_4| = \binom{n-1}{2} 2(n-1)^{n-4} = (n-2)p_{n-2}$. Therefore, $p_{n,1}^n - p_{n,1}^{n-1} = p_{n-2}$ for any $n \geq 2$. ■

To enumerate preference sets in $\mathcal{P}^l_{n;\leq s;k}$ with $k \leq l \leq s$, we need the following lemma.

Lemma 6.2. *Suppose $k \geq 1$, $k \leq l \leq s-1$ and $s \leq n$. Let $\tilde{\mathcal{P}}^l_{n;\leq s;k} = \{\alpha \in \mathcal{P}^l_{n;\leq s;k} \mid m_\alpha < l\}$. Then $|\tilde{\mathcal{P}}^l_{n;\leq s;k}| = \sum\limits_{i=k-1}^{l-2} \binom{n-1}{i+1-k} p_{i+1-k,i;\leq i} p^{l-i-1}_{n+k-i-1;\leq s-i-1}$.*

Proof. For any $\alpha = (a_1, \ldots, a_n) \in \tilde{\mathcal{P}}^l_{n;\leq s;k}$, we suppose that the last empty parking space is $i+1$, then $k-1 \leq i \leq l-2$. Let $S = \{j \mid 1 \leq a_j \leq i\}$ and α_S be a subsequence of α determined by the subscripts in S, then $|S| = i+1-k$ and $\alpha_S \in \mathcal{P}_{i+1-k,i;\leq i}$. Let $T = [n] \setminus S$ and α_T be a subsequence of α determined by the subscripts in T, then $|T| = n+k-i-1$. Suppose $\alpha_T = (b_1, \ldots, b_{n+k-i-1})$, since $i+2 \leq a_j \leq s$ for any $j \in T$, we have $(b_1 - i - 1, \ldots, b_{n+k-i-1} - i - 1) \in \mathcal{P}^{l-i-1}_{n+k-i-1;\leq s-i-1}$.

There are $\binom{n-1}{i+1-k}$ ways to choose $i+1-k$ numbers from $[2,n]$ for the elements in S since $1 \notin S$. There are $p_{i+1-k,i;\leq i}$ and $p^{l-i-1}_{n+k-i-1;\leq s-i-1}$ possibilities for the preference set α_S and the parking function α_T, respectively. Hence, we have

$$|\tilde{\mathcal{P}}^l_{n;\leq s;k}| = \sum_{i=k-1}^{l-2} \binom{n-1}{i+1-k} p_{i+1-k,i;\leq i} p^{l-i-1}_{n+k-i-1;\leq s-i-1}.$$

■

Example 6.1. *Take $n = 7$, $s = 6$, $l = 4$ and $k = 2$. By the computer search, we have $|\tilde{\mathcal{P}}^4_{7;\leq 6;2}| = 6265$. By the data in Appendix, we find $p_{0,1;\leq 1} = 1$, $p^2_{7;\leq 4} = 3361$, $p_{1,2;\leq 2} = 2$ and $p^1_{6;\leq 3} = 242$. Hence, it is easy to check that $|\tilde{\mathcal{P}}^4_{7;\leq 6;2}| = \sum\limits_{i=1}^{2} \binom{6}{i-1} p_{i-1,i;\leq i} p^{3-i}_{8-i;\leq 5-i}$.*

Theorem 6.6. *Let n, s, k, l be integers. Suppose $k \geq 1$, $k \leq l \leq s-1$ and $s \leq n$. Then*

$$\begin{aligned} p^{l+1}_{n;\leq s;k} - p^l_{n;\leq s;k} &= \sum_{i=k-1}^{l-2} \binom{n-1}{i-k+1} p_{i-k+1,i;\leq i} \left[p^{l-i}_{n+k-i-1;\leq s-i-1} - p^{l-i-1}_{n+k-i-1;\leq s-i-1}\right] \\ &\quad + \binom{n-1}{l-k} p_{l-k,l-1;\leq l-1} \left[p^1_{n+k-l;\leq s-l} - p^{s-l}_{n+k-l;\leq s-l;1}\right]. \end{aligned}$$

Proof. Let $\mathcal{A}'_1$ and $\mathcal{C}'_1$ defined as that in Lemma 5.2, $\mathcal{A}'_2$ and $\mathcal{C}'_2$ defined as that in Lemma 5.3, $\mathcal{A}'_3$ defined as that in Lemma 5.4, $\tilde{\mathcal{P}}^{l+1}_{n,\leq s,k}$ and $\tilde{\mathcal{P}}^l_{n,\leq s,k}$ defined as that in Lemma 6.2. Obviously, $\mathcal{P}^l_{n;\leq s;k} = \bigcup\limits_{i=1}^{3} \mathcal{A}'_i \cup \tilde{\mathcal{P}}^l_{n;\leq s;k}$ and $\mathcal{P}^{l+1}_{n;\leq s;k} = \bigcup\limits_{i=1}^{2} \mathcal{C}'_i \cup \tilde{\mathcal{P}}^{l+1}_{n;\leq s;k}$. Hence,

$$|\mathcal{P}^{l+1}_{n;\leq s;k}| - |\mathcal{P}^l_{n;\leq s;k}| = |\tilde{\mathcal{P}}^{l+1}_{n;\leq s;k}| - |\tilde{\mathcal{P}}^l_{n;\leq s;k}| - |\mathcal{A}'_3|.$$

By Lemmas 5.5 and 6.2, we have

$$
\begin{aligned}
& p^{l+1}_{n;\leq s;k} - p^{l}_{n;\leq s;k} \\
= & \sum_{i=k-1}^{l-1} \binom{n-1}{i+1-k} p_{i+1-k,i;\leq i} p^{l-i}_{n+k-i-1;\leq s-i-1} - \sum_{i=k-1}^{l-2} \binom{n-1}{i+1-k} p_{i+1-k,i;\leq i} p^{l-i-1}_{n+k-i-1;\leq s-i-1} \\
& - \binom{n-1}{l-k} p_{l-k,l-1;\leq l-1} p^{s-l}_{n+k-l;\leq s-l;1} \\
= & \sum_{i=k-1}^{l-2} \binom{n-1}{i-k+1} p_{i-k+1,i;\leq i} \left[p^{l-i}_{n+k-i-1;\leq s-i-1} - p^{l-i-1}_{n+k-i-1;\leq s-i-1} \right] \\
& + \binom{n-1}{l-k} p_{l-k,l-1;\leq l-1} \left[p^{1}_{n+k-l;\leq s-l} - p^{s-l}_{n+k-l;\leq s-l;1} \right].
\end{aligned}
$$

■

Example 6.2. *Take* $n = 7, s = 6$, $l = 4$ *and* $k = 1$. *By the data in Appendix, we find* $p^5_{7;\leq 6;1} = 19042$, $p^4_{7;\leq 6;1} = 18602$. *Hence,* $p^5_{7;\leq 6;1} - p^4_{7;\leq 6;1} = 440$. *On the other hand, we have* $p_{0,0;\leq 0} = 1$, $p^4_{7;\leq 5} = 9351$, $p^3_{7,\leq 5} = 10026$, $p_{1,1,\leq 1} = 1$, $p^3_{6,\leq 4} = 701$, $p^2_{6;\leq 4} = 776$, $p_{2,2;\leq 2} = 3$, $p^2_{5;\leq 3} = 65$, $p^1_{5,\leq 3} = 80$, $p_{3,3,\leq 3} = 16$, $p^1_{4,\leq 2} = 8$, $p^2_{4;\leq 2;1} = 1$. *It is easy to check that* $p^5_{7;\leq 6;1} - p^4_{7;\leq 6;1} = \binom{6}{3} p_{3,3;\leq 3} \left[p^1_{4;\leq 2} - p^2_{4;\leq 2;1} \right] + \sum_{i=0}^{2} \binom{6}{i} p_{i,i;\leq i} \left[p^{4-i}_{7-i;\leq 5-i} - p^{3-i}_{7-i;\leq 5-i} \right]$.

7 The generating function

In this section, we will study some generating functions for some sequences given in the previous sections. First, we need the following three lemmas [11].

Lemma 7.1. [11] *Suppose that* $k \geq 0$. *Let* $Q_k(x) = \sum_{n \geq 0} \frac{p_{n,n+k;\leq n+k}}{n!} x^n$, *then* $Q_k(x) = [P(x)]^{k+1}$.

Lemma 7.2. [11] *Suppose that* $k \geq 0$. *Let* $R_k(x) = \sum_{n \geq k} \frac{p_{n;\leq n-k}}{n!} x^n$ *and* $R(x,y) = \sum_{k \geq 0} R_k(x) y^k$, *then* $R_k(x) = P(x) \sum_{i=0}^{k} \frac{(-1)^i (k+1-i)^i}{i!} x^i - \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i$ *and* $R(x,y) = \frac{P(x)-y}{e^{xy}-y}$.

Lemma 7.3. [11] *Let* $F(x,y,z) = \sum_{k \geq 0} \sum_{s \geq k} \sum_{n \geq s+1} \frac{p^{n-s}_{n;\leq n-k}}{(n-1)!} x^n y^s z^k$, *then*

$$
F(x,y,z) = \frac{x}{e^{xyz} - z} \left[\frac{P(xy)(P(x) - yP(xy))}{1-y} - \frac{zP(xyz)[P(x) - yzP(xyz)]}{1-yz} \right].
$$

Now, for any $k \geq 1$ and $s \geq 0$, we define a generating function $D_{k,s}(x) = \sum_{n \geq s+k} \frac{p_{n;\leq n-s;k}}{n!} x^n$.

Theorem 7.1. *Suppose $s \geq 0$ and $k \geq 1$. Let $D_{k,s}(x)$ be the generating function for $p_{n;\leq n-s;k}$, then*

$$D_{k,s}(x) = [P(x)]^{k+1} \sum_{i=0}^{k+s} \frac{(-1)^i (k+s+1-i)^i}{i!} x^i - [P(x)]^k \sum_{i=0}^{k+s-1} \frac{(-1)^i (k+s-i)^i}{i!} x^i.$$

Proof. By Lemma 2.1, we have

$$\begin{aligned}
\sum_{n \geq s+k} p_{n;\leq n-s;k} \frac{x^n}{n!} &= \sum_{n \geq s+k} \sum_{i=1}^{n-s-k} \binom{n}{s+i+k} p_{n-s-i-k,n-s-i-1;\leq n-s-i-1} p_{s+i+k;\leq i} \frac{x^n}{n!} \\
&= \sum_{n \geq 0} \sum_{i=0}^{n} \frac{p_{n-i,n+k-i-1;\leq n+k-i-1}}{(n-i)!} \frac{p_{s+i+k;\leq i}}{(s+i+k)!} x^{n+s+k},
\end{aligned}$$

hence,

$$\begin{aligned}
D_{k,s}(x) &= Q_{k-1}(x) R_{s+k}(x) \\
&= [P(x)]^{k+1} \sum_{i=0}^{k+s} \frac{(-1)^i (k+s+1-i)^i}{i!} x^i - [P(x)]^k \sum_{i=0}^{k+s-1} \frac{(-1)^i (k+s-i)^i}{i!} x^i.
\end{aligned}$$

■

Corollary 7.1. *Let $D_k(x,y) = \sum_{s \geq 0} D_{k,s}(x) y^s$ for any $k \geq 1$, then*

$$D_k(x,y) = \left[\frac{P(x)}{y}\right]^k \left[R(x,y) - \sum_{s=0}^{k-1} R_s(x) y^s\right].$$

Furthermore, let $D(x,y,z) = \sum_{k \geq 1} D_k(x,y) z^k$, then

$$D(x,y,z) = \frac{zP(x)}{y - zP(x)} \left[\frac{P(x) - y}{e^{xy} - y} - \frac{(1-z)P(x)}{e^{xzP(x)} - zP(x)}\right].$$

Proof. By Theorem 7.1, we have

$$\begin{aligned}
D_k(x,y) &= \sum_{s \geq 0} D_{k,s}(x) y^s \\
&= \sum_{s \geq 0} [P(x)]^k R_{s+k}(x) y^s \\
&= \left[\frac{P(x)}{y}\right]^k \left[R(x,y) - \sum_{s=0}^{k-1} R_s(x) y^s\right].
\end{aligned}$$

Hence,

$$\begin{aligned} D(x,y,z) &= \sum_{k\geq 1}\left[\frac{P(x)}{y}\right]^k\left[R(x,y)-\sum_{s=0}^{k-1}R_s(x)y^s\right]z^k \\ &= \sum_{k\geq 0}\left[\frac{zP(x)}{y}\right]^{k+1}R(x,y)-\sum_{k\geq 0}\left[\frac{zP(x)}{y}\right]^{k+1}\sum_{s=0}^{k}R_s(x)y^s \\ &= \frac{zP(x)}{y-zP(x)}[R(x,y)-R(x,zP(x)] \\ &= \frac{zP(x)}{y-zP(x)}\left[\frac{P(x)-y}{e^{xy}-y}-\frac{(1-z)P(x)}{e^{xzP(x)}-zP(x)}\right]. \end{aligned}$$

■

For any $k \geq 1$, we define a generating function $I_k(x) = \sum_{n\geq 1}\frac{p^1_{n,n+k;\leq n+k}}{(n-1)!}x^n$ and let $I(x,y) = \sum_{k\geq 0} I_k(x)y^k$.

Lemma 7.4. *Suppose that $k \geq 0$. Let $I_k(x)$ be the generating function for $p^1_{n,n+k;\leq n+k}$, then $I_k(x)$ satisfies the following recurrence relation*

$$I_k(x) = I_{k-1}(x)P(x)$$

for any $k \geq 1$, with the initial condition $I_0(x) = x[P(x)]^2$. Equivalently,

$$I_k(x) = x[P(x)]^{k+2}.$$

Let $I(x,y) = \sum_{k\geq 0} I_k(x)y^k$, then

$$I(x,y) = \frac{x[P(x)^2]}{1-yP(x)}.$$

Proof. When $k = 0$, it is known that $I_0(x) = x[P(x)]^2$. When $k \geq 1$, Theorem 4.2 implies that

$$\sum_{n\geq 1}\frac{p^1_{n,n+k;\leq n+k}}{(n-1)!}x^n = \sum_{n\geq 1}\sum_{i=1}^{n}\frac{p^1_{i,i+k-1;\leq i+k-1}}{(i-1)!}\frac{p_{n-i}}{(n-i)!}x^n.$$

Hence,

$$I_k(x) = I_{k-1}(x)P(x) \quad \text{and} \quad I_k(x) = x[P(x)]^{k+2}.$$

It is easy to obtain that

$$I(x,y) = \frac{x[P(x)]^2}{1-yP(x)}$$

$\blacksquare$

For any $l \geq 0$ and $k \geq 0$, we define a generating function $H_{l,k}(x) = \sum_{n\geq l+1} \frac{p^{n+k-l}_{n,n+k;\leq n+k}}{(n-1)!}x^n$

Theorem 7.2. *Suppose $l \geq 0$ and $k \geq 0$. Let $H_{l,k}(x)$ be the generating function for $p^{n+k-l}_{n,n+k;\leq n+k}$, then $H_{l,k}(x)$ satisfies the following recurrence relation*

$$H_{l,k}(x) = H_{l-1,k}(x) - \frac{p^{1}_{l,l+k;\leq l+k}}{(l-1)!}x^l + \frac{p_l}{l!}x^{l+1}[P(x)]^{k+1}$$

with the initial conditions $H_{0,k}(x) = x[P(x)]^{k+1}$.

Proof. When $l = 0$, Lemma 4.1 tells us that

$$p^{n+k}_{n,n+k;\leq n+k} = p_{n-1,n+k-1;\leq n+k-1}$$

for any $k \geq 0$ and $n \geq 1$. Hence,

$$\begin{aligned} H_{0,k}(x) &= \sum_{n\geq 1} \frac{p^{n+k}_{n,n+k;\leq n+k}}{(n-1)!}x^n \\ &= \sum_{n\geq 1} \frac{p_{n-1,n+k-1;\leq n+k-1}}{(n-1)!}x^n \\ &= x\sum_{n\geq 0} \frac{p_{n,n+k;\leq n+k}}{n!}x^n \\ &= xQ_k(x) \\ &= x[P(x)]^{k+1}. \end{aligned}$$

Given $l \geq 1$, by Theorem 4.3, we have

$$p^{n+k-l}_{n,n+k;\leq n+k} - p^{n+k-l+1}_{n,n+k;\leq n+k} = \binom{n-1}{l} p_{n-l-1,n+k-l-1;\leq n+k-l-1}p_l.$$

This implies that

$$\begin{aligned} H_{l,k}(x) &= H_{l-1,k}(x) - \frac{p^{k+1}_{l,l+k;\leq l+k}}{(l-1)!}x^l + x^{l+1}\frac{p_l}{l!}Q_k(x) \\ &= H_{l-1,k}(x) - \frac{p^{1}_{l,l+k;\leq l+k}}{(l-1)!}x^l + \frac{p_l}{l!}x^{l+1}[P(x)]^{k+1}. \end{aligned}$$

$\blacksquare$

Corollary 7.2. *Let* $H_k(x,y)=\sum_{l\geq 0} H_{l,k}(x)y^l$ *, then* $H_k(x)$ *satisfies the following equation*

$$H_k(x,y)=yH_k(x,y)-xy[P(xy)]^{k+2}+x[P(x)]^{k+1}P(xy).$$

Equivalently,

$$H_k(x,y)=\frac{xP(xy)\left\{[P(x)]^{k+1}-y[P(xy)]^{k+1}\right\}}{1-y}.$$

Let $H(x,y,z)=\sum_{k\geq 0} H_k(x,y)z^k$*, then*

$$H(x,y,z)=\frac{xP(xy)}{1-y}\left[\frac{P(x)}{1-zP(x)}-\frac{yP(xy)}{1-zP(xy)}\right].$$

Proof. By Theorem 7.2, we have

$$\sum_{l\geq 1} H_{l,k}(x)y^l=\sum_{l\geq 1} H_{l-1,k}(x)y^l-\sum_{l\geq 1}\frac{p^1_{l,l+k;\leq l+k}}{(l-1)!}(xy)^l+x[P(x)]^{k+1}\sum_{l\geq 1}\frac{p_l}{l!}(xy)^l.$$

Therefore, $H_k(x,y)=yH_k(x,y)-xy[P(xy)]^{k+2}+x[P(x)]^{k+1}P(xy)$. Equivalently, we have

$$H_k(x,y)=\frac{xP(xy)\left\{[P(x)]^{k+1}-y[P(xy)]^{k+1}\right\}}{1-y}.$$

Furthermore,

$$H(x,y,z)=\frac{xP(xy)}{1-y}\left[\frac{P(x)}{1-zP(x)}-\frac{yP(xy)}{1-zP(xy)}\right].$$

■

Let $k\geq 1$ and $s\geq 0$. Define a generating function $M_{s,k}(x)=\sum_{n\geq s+k+2}\frac{p^1_{n;\leq n-s;k}}{(n-1)!}x^n$.

Lemma 7.5. *Suppose that* $k\geq 1$ *and* $s\geq 0$*. Let* $M_{s,k}(x)$ *be the generating function for* $p^1_{n;\leq n-s;k}$*, then*

$$M_{s,k}(x)=R_{s+k}(x)I_{k-1}(x).$$

Let $M_k(x,y)=\sum_{s\geq 0} M_{s,k}(x)y^s$ *and* $M(x,y,z)=\sum_{k\geq 1} M_k(x,y)z^k$*, then*

$$M_k(x,y)=I_{k-1}(x)y^{-k}\left[R(x,y)-\sum_{s=0}^{k-1}R_s(x)y^s\right]$$

and

$$M(x,y,z)=\frac{xz[P(x)]^2}{y-zP(x)}\left[\frac{P(x)-y}{e^{xy}-y}-\frac{(1-z)P(x)}{e^{xzP(x)}-zP(x)}\right].$$

Proof. For any $k \geq 1$ and $s \geq 0$, Theorem 6.2 implies that

$$p^1_{n;\leq n-s;k} = \sum_{i=1}^{n-s-k-1} \binom{n-1}{s+k+i} p_{s+k+i;\leq i} p^1_{n-s-k-i,n-s-i-1;\leq n-s-i-1},$$

hence,

$$M_{s,k}(x) \;=\; R_{s+k}(x) I_{k-1}(x).$$

So,

$$\begin{aligned} M_k(x,y) &= \sum_{s\geq 0} R_{s+k}(x) I_{k-1}(x) y^s \\ &= I_{k-1}(x) y^{-k} \left[R(x,y) - \sum_{s=0}^{k-1} R_s(x) y^s \right] \end{aligned}$$

Furthermore, we have

$$\begin{aligned} M(x,y,z) &= \sum_{k\geq 1} I_{k-1}(x) y^{-k} [R(x,y) - \sum_{s=0}^{k-1} R_s(x) y^s] z^k \\ &= \frac{xz[P(x)]^2}{y - P(x)z} \left[R(x,y) - R(x, zP(x)) \right] \\ &= \frac{xz[P(x)]^2}{y - zP(x)} \left[\frac{P(x) - y}{e^{xy} - y} - \frac{(1-z)P(x)}{e^{xzP(x)} - zP(x)} \right]. \end{aligned}$$

■

Define a generating function

$$W(x,y,z,v) = \sum_{k\geq 1} \sum_{s\geq 0} \sum_{l\geq s} \sum_{n\geq k+l} \frac{p^{n-l}_{n;\leq n-s;k}}{(n-1)!} x^n y^l z^s v^k.$$

Theorem 7.3.

$$\begin{aligned} W(x,y,z,v) \;=\; & \frac{xyvP(xy)}{y-1} \left\{ \left[\frac{yP(xy)}{yz - vP(xy)} - \frac{P(x)}{yz - vP(x)} \right] R(xy,z) \right. \\ & \left. + \frac{P(x)}{yz - vP(x)} R(xy, \frac{v}{y} P(x)) - \frac{yP(xy)}{yz - vP(xy)} R(xy, \frac{v}{y} P(xy)) \right\} \\ & + \frac{vP(x)}{yz - vP(x)} \left[F(x,y,z) - F(x,y,\frac{v}{y} P(x)) \right] \end{aligned}$$

Proof. First, let $W_{k,s,l}(x) = \sum_{n\geq k+l} \frac{p^{n-l}_{n;\leq n-s;k}}{(n-1)!} x^n$. When $l = s = 0$, Theorems 6.5 and 6.3 imply that

$$p^n_{n;k} = \begin{cases} p^2_n = p^1_n - p_{n-1} & if \quad k = 1 \\ p^1_{n;k-1} & if \quad k \geq 2 \end{cases}.$$

This tells us that

$$W_{k,0,0}(x)=\begin{cases} xP(x)[P(x)-1] & if \quad k=1\\ M_{0,k-1}(x) & if \quad k\geq 2\end{cases}.$$

When $l=s\geq 1$, by Theorem 6.4, we have $W_{k,s,s}(x)=xD_{k,s-1}(x)$.

For any $l\geq s+1$, by Theorem 6.6, it follows that

$$\begin{aligned} p^{n-l+1}_{n;\leq n-s;k}-p^{n-l}_{n;\leq n-s;k} &= \sum_{i=0}^{n-k-l-1}\binom{n-1}{i}p_{i,i+k-1;\leq i+k-1}\left[p^{n-k-l+1-i}_{n-i;\leq n-s-k-i}-p^{n-k-l-i}_{n-i;\leq n-s-k-i}\right]\\ &\quad+\binom{n-1}{n-k-l}p_{n-k-l,n-l-1;\leq n-l-1}\left[p^{1}_{l+k;\leq l-s}-p_{l+k-1;\leq l-s;1}\right]\\ &= \sum_{i=0}^{n-k-l}\binom{n-1}{i}p_{i,i+k-1;\leq i+k-1}\left[p^{n-k-l+1-i}_{n-i;\leq n-s-k-i}-p^{n-k-l-i}_{n-i;\leq n-s-k-i}\right]\\ &\quad-\binom{n-1}{n-k-l}p_{n-k-l,n-l-1;\leq n-l-1}p_{l+k-1;\leq l-s;1}.\end{aligned}$$

Let $F_{k,s}(x)=\sum\limits_{n\geq s+1}\frac{p^{n-s}_{n;\leq n-k}}{(n-1)!}x^n$, then

$$\begin{aligned} &W_{k,s,l-1}(x)-\frac{p^{k}_{k+l-1;\leq k+l-1-s;k}}{(k+l-2)!}x^{k+l-1}-W_{k,s,l}(x)\\ =\ &\left[-\frac{p_{l+k-1;\leq l-s;1}}{(k+l-1)!}x^{k+l}+F_{s+k,k+l-1}(x)-F_{s+k,k+l}(x)\right]Q_{k-1}(x).\end{aligned}$$

Let $W_{k,s}(x,y)=\sum\limits_{l\geq s}\sum\limits_{n\geq k+l}\frac{p^{n-l}_{n;\leq n-s;k}}{(n-1)!}x^ny^l=\sum\limits_{l\geq s}W_{k,s,l}(x)y^l$ and $F_k(x,y)=\sum\limits_{s\geq k}F_{k,s}(x)y^s$, then

$$\begin{aligned} &yW_{k,s}(x,y)-y^{-k+1}M_{s,k}(xy)-W_{k,s}(x,y)+W_{k,s,s}(x)y^s\\ =\ &\left[-D_{1,s+k-1}(xy)y^{-k+1}x+F_{s+k}(x,y)y^{-k}(y-1)+F_{s+k,s+k}(x)y^s\right]Q_{k-1}(x).\end{aligned}$$

Furthermore, note that $F(x,y,z)=\sum\limits_{k\geq 0}F_k(x,y)z^k$ and let $W_k(x,y,z)=\sum\limits_{s\geq 0}\sum\limits_{l\geq s}\sum\limits_{n\geq k+l}\frac{p^{n-l}_{n;\leq n-s;k}}{(n-1)!}x^ny^lz^s=\sum\limits_{s\geq 0}W_{k,s}(x,y)z^s$, then

$$\begin{aligned} &(y-1)W_k(x,y,z)-y^{-k+1}M_k(xy,z)+W_{k,0,0}(x)+xyzD_k(x,yz)\\ =\ &-\left[D_1(xy,z)-\sum_{s=0}^{k-2}D_{1,s}(xy)z^s\right](yz)^{-k+1}xQ_{k-1}(x)\\ &+\left[F(x,y,z)-\sum_{s=0}^{k-1}F_s(x,y)z^s\right](yz)^{-k}(y-1)Q_{k-1}(x)\\ &+x(yz)^{-k+1}\left[R(x,yz)-\sum_{s=0}^{k-2}R_s(x)(yz)^s\right]Q_{k-1}(x)\\ &-x^k\sum_{s\geq k-1}\frac{p_{s;\leq 0}}{s!}(xyz)^{s-k+1}Q_{k-1}(x).\end{aligned}$$

Since $W(x,y,z,v)=\sum_{k\geq 1}\sum_{s\geq 0}\sum_{l\geq s}\sum_{n\geq k+l}\frac{p^{n-l}_{n;\leq n-s;k}}{(n-1)!}x^n y^l z^s v^k=\sum_{k\geq 1}W_k(x,y,z)v^k$, we have

$$\begin{aligned}
& (y-1)W(x,y,z,v)-yM(xy,z,\frac{v}{y})+vxP(x)[R(x,vP(x))-1]+xyzD(x,yz,v)\\
= & -\frac{xyzvP(x)}{yz-vP(x)}D_1(xy,z)+\frac{xv^2[P(x)]^2}{yz-vP(x)}D_1(xy,\frac{v}{y}P(x))\\
& +\frac{(y-1)vP(x)}{yz-vP(x)}\left[F(x,y,z)-F(x,y,\frac{v}{y}P(x))\right]\\
& +\frac{xyzvP(x)}{yz-vP(x)}R(x,yz)-\frac{xv^2[P(x)]^2}{yz-vP(x)}R(x,vP(x))-xvP(x)
\end{aligned}$$

Therefore,

$$\begin{aligned}
W(x,y,z,v) = & \frac{xyvP(xy)}{y-1}\left[\frac{yP(xy)}{yz-vP(xy)}-\frac{P(x)}{yz-vP(x)}\right]R(xy,z)\\
& +\frac{xyvP(xy)}{y-1}\left[\frac{P(x)}{yz-vP(x)}R(xy,\frac{v}{y}P(x))-\frac{yP(xy)}{yz-vP(xy)}R(xy,\frac{v}{y}P(xy))\right]\\
& +\frac{vP(x)}{yz-vP(x)}\left[F(x,y,z)-F(x,y,\frac{v}{y}P(x))\right]
\end{aligned}$$

■

8 Appendix

For convenience to check the equations given in the previous sections, by the computer search, for $n\leq 7$, we obtain the number of k-flaw preference sets $\alpha=(a_1,\ldots,a_n)$ of length n satisfying $a_1=l$ and $a_i\leq s$ for all $i\in[n]$ and list them in the following tables. Note that $p^l_{n;\leq s;k}=0$ if $l>s$ or $n\leq k$. In Table 5, we give the values of $p^l_{n,n+k;\leq n+k}$ for any $n\leq 5$ and $k\leq 3$.

$k = 0$	$l = 1$	2	3	4	5	6	7	8	$p_{n,\leq s}$
$(n, s) = (1, 1)$	1	0							1
$(2, 1)$	1	0							1
$(2, 2)$	2	1	0						3
$(3, 1)$	1	0							1
$(3, 2)$	4	3	0						7
$(3, 3)$	8	5	3	0					16
$(4, 1)$	1	0							1
$(4, 2)$	8	7	0						15
$(4, 3)$	26	19	16	0					61
$(4, 4)$	50	34	25	16	0				125
$(5, 1)$	1	0							1
$(5, 2)$	16	15	0						31
$(5, 3)$	80	65	61	0					206
$(5, 4)$	232	171	143	125	0				671
$(5, 5)$	432	307	243	189	125	0			1296
$(6, 1)$	1	0							1
$(6, 2)$	32	31							63
$(6, 3)$	242	211	206	0					659
$(6, 4)$	982	776	701	671					3130
$(6, 5)$	2642	1971	1666	1456	1296	0			9031
$(6, 6)$	4802	3506	2881	2401	1921	1296	0		16807
$(7, 1)$	1	0							1
$(7, 2)$	64	63	0						127
$(7, 3)$	728	665	659	0					2052
$(7, 4)$	4020	3361	3175	3130	0				13686
$(7, 5)$	14392	11262	10026	9351	9031	0			54062
$(7, 6)$	36724	27693	23667	20922	18682	16807	0		144495
$(7, 7)$	65536	48729	40953	35328	30208	24583	16807	0	262144

Table.1. $p^l_{n;\leq s}$ for $1 \leq n \leq 7$

$k=1$	$l=1$	2	3	4	5	6	7	8	$p_{n,\leq s,1}$
$(n,s)=(2,1)$	0	0							0
$(2,2)$	0	1	0						1
$(3,1)$	0	0							0
$(3,2)$	0	1	0						1
$(3,3)$	1	4	5	0					10
$(4,1)$	0	0							0
$(4,2)$	0	1	0						1
$(4,3)$	1	8	10	0					19
$(4,4)$	13	29	31	34	0				107
$(5,1)$	0	0							0
$(5,2)$	0	1	0						1
$(5,3)$	1	616	19	0					36
$(5,4)$	23	84	97	107	0				311
$(5,5)$	165	290	293	291	307	0			1346
$(6,1)$	0	0							0
$(6,2)$	0	1	0						1
$(6,3)$	1	32	36	0					69
$(6,4)$	41	247	291	311	0				890
$(6,5)$	436	1107	1206	1266	1346	0			5361
$(6,6)$	2341	3637	3591	3461	3381	3506	0		19917
$(7,1)$	0	0							0
$(7,2)$	0	1	0						1
$(7,3)$	1	64	69	0					134
$(7,4)$	75	734	857	890	0				2556
$(7,5)$	1151	4281	4858	5161	5361	0			20812
$(7,6)$	8402	17433	18329	18602	19042	19917	0		101725
$(7,7)$	37883	54690	53435	51008	48808	47433	48729	0	341986

Table.2. $p^{l}_{n;\leq s;1}$ for $2 \leq n \leq 7$

$k=2$	$l=1$	2	3	4	5	6	7	8	$p_{n,\leq s,2}$
$(n,s)=(3,1)$	0	0							0
$(3,2)$	0	0	0						0
$(3,3)$	0	0	1	0					1
$(4,1)$	0	0							0
$(4,2)$	0	0	0						0
$(4,3)$	0	0	1	0					1
$(4,4)$	1	1	8	13	0				23
$(5,1)$	0	0							0
$(5,2)$	0	0	0						0
$(5,3)$	0	0	1	0					1
$(5,4)$	1	1	16	23	0				41
$(5,5)$	27	27	88	129	165	0			436
$(6,1)$	0	0							0
$(6,2)$	0	0	0						0
$(6,3)$	0	0	1	0					1
$(6,4)$	1	1	32	41	0				75
$(6,5)$	46	46	252	371	436	0			1151
$(6,6)$	581	581	1252	1656	1991	2341	0		8402
$(7,1)$	0	0							0
$(7,2)$	0	0	0						0
$(7,3)$	0	0	1	0					1
$(7,4)$	1	1	64	75	0				141
$(7,5)$	81	81	740	1049	1151	0			3102
$(7,6)$	1442	1442	4572	6385	7627	8402	0		29870
$(7,7)$	12373	12373	21404	26326	29938	33563	37883	0	173860

Table.3. $p^{l}_{n;\leq s;2}$ for $3 \leq n \leq 7$

$k = 3$	$l = 1$	2	3	4	5	6	7	8	$p_{n,\leq s,3}$
$(n, s) = (4, 1)$	0	0							0
$(4, 2)$	0	0	0						0
$(4, 3)$	0	0	0	0					0
$(4, 4)$	0	0	0	1	0				1
$(5, 1)$	0	0							0
$(5, 2)$	0	0	0						0
$(5, 3)$	0	0	0	0					0
$(5, 4)$	0	0	0	1	0				1
$(5, 5)$	1	1	1	16	27	0			46
$(6, 1)$	0	0							0
$(6, 2)$	0	0	0						0
$(6, 3)$	0	0	0	0					0
$(6, 4)$	0	0	0	1	0				1
$(6, 5)$	1	1	1	32	46	0			81
$(6, 6)$	51	51	51	257	451	581	0		1442
$(7, 1)$	0	0							0
$(7, 2)$	0	0	0						0
$(7, 3)$	0	0	0	0					0
$(7, 4)$	0	0	0	1	0				1
$(7, 5)$	1	1	1	64	81	0			148
$(7, 6)$	87	87	87	746	1241	1442	0		3690
$(7, 7)$	1763	1763	1763	4893	7942	10573	12373	0	41070

Table.4. $p^{l}_{n;\leq s;3}$ for $4 \leq n \leq 7$

	$l = 1$	2	3	4	5	6	7	8	$p_{n,n+k;\leq n+k}$
$(n, k) = (1, 0)$	1								1
$(1, 1)$	1	1							2
$(1, 2)$	1	1	1						3
$(1, 3)$	1	1	1	1					4
$(2, 0)$	2	1							3
$(2, 1)$	3	3	2						8
$(2, 2)$	4	4	4	3					15
$(2, 3)$	5	5	5	5	4				24
$(3, 0)$	8	5	3						16
$(3, 1)$	15	15	12	8					50
$(3, 2)$	24	24	24	21	15				108
$(3, 3)$	35	35	35	35	32	24			196
$(4, 0)$	50	34	25	16					125
$(4, 1)$	108	108	92	74	50				432
$(4, 2)$	196	196	196	180	153	108			1029
$(4, 3)$	320	320	320	320	304	268	196		2048
$(5, 0)$	432	307	243	189	125				1296
$(5, 1)$	1029	1029	904	776	632	432			4802
$(5, 2)$	2048	2048	2048	1923	1731	1461	1029		12288
$(5, 3)$	3645	3645	3645	3645	3520	3264	2832	2048	26244

Table.5. $p^{l}_{n,n+k;\leq n+k}$ for $1 \leq n \leq 5$ and $k \leq 3$

References

[1] Peter J Cameron, Daniel Johannsen, Thomas Prellberg, Pascal Schweitzer, Couting Defective Parking Functions, arXiv:0803.0302v1, 3 Mar, 2008

[2] R. Cori, D. Rossin, B. Salvy, Polynomial ideals for sandpiles and their Grobner bases. *Theoretical Computer Science* **276** (2002), no. 1-2, 1-15.

[3] Sen-Peng Eu,Tung-Shan Fu,Chun-Ju Lai,On the enumeration of parking functions by leading terms, *Adv. in Appl. Math.* 35 (2005) 392-406

[4] Sen-Peng Eu, Tung-Shan Fu, Yeong-NanYeh, Refined Chung-Feller theorems for lattice paths, *J. Combin. Theory, Ser. A* 112 (2005) 143-162.

[5] Sen-Peng Eu, Shu-Chung Liu, Yeong-NanYeh, Taylor expansions for Catalan and Motzkin numbers, *Adv. in Appl. Math.* 29 (2002) 345-357.

[6] J. Françon, Acyclic and parking functions, *J. Combin. Theory Ser. A* 18 (1975) 27-35.

[7] D. Foata, J. Riordan, Mappings of acyclic and parking functions, *Aequationes Math.* 10 (1974) 10-22.

[8] J.D. Gilbey, L.H. Kalikow, Parking functions, valet functions and priority queues, *Discrete Math.* 197/198 (1999) 351-373.

[9] A. Henderson. The twenty-seven lines upon the cubic surface, Cambridge University Press (1911)

[10] Po-Yi Huang, Jun Ma, Yeong-Nan Yeh, Ordered k-flaw Preference sets, submitted.

[11] Po-Yi Huang, Jun Ma, Chun-Chen Yeh, Some enumerations for parking functions, submitted.

[12] J.P.S. Kung, C.H. Yan, Gončarove polynomials and parking functions, *J. Combin. Theory Ser. A* 102 (2003) 16C37.

[13] J. Pitman, R. Stanley, A polytope related to empirical distributions, plane trees, parking functions, and the associahedron, *Discrete Comput. Geom.* 27 (4) (2002) 603-634.

[14] Postnikov, A. and Shapiro, B. Trees, Parking Functions, Syzygies, and Deformatioins of Monomial Ideals. *Transactions of the American Mathematical Society* **356** (2004).

[15] J.Riordan, Ballots and trees, *J.Combin. Theory* 6 (1969) 408-411.

[16] M.P. Schtzenberger, On an enumeration problem, *J. Combin. Theory* 4 (1968) 219-221.

[17] R.P. Stanley, Hyperplane arrangements, interval orders and trees, *Proc. Natl. Acad. Sci.* 93 (1996) 2620-2625.

[18] R.P. Stanley, Parking functions and non-crossing partitions, in: The Wilf Festschrift, *Electron. J. Combin.* 4 (1997) R20.

[19] C.H. Yan, Generalized tree inversions and k-parking functions, *J. Combin. Theory Ser. A* 79 (1997) 268-280.

[20] C.H. Yan, On the enumeration of generalized parking functions, *Congr. Numer.* 147 (2000) 201-209.

[21] C.H. Yan, Generalized parking functions, tree inversions and multicolored graphs, *Adv. in Appl. Math.* 27 (2001) 641-670.